

INTERNATIONAL SCIENTIFIC
PERIODICAL JOURNAL



THE EUROPEAN ASSOCIATION OF PEDAGOGUES AND PSYCHOLOGISTS "SCIENCE"

"THE UNITY OF SCIENCE"

MONOGRAPH

Buklina S.B.
COGNITIVE DISORDERS
IN PATIENTS WITH LESIONS OF DEEP CEREBRAL
STRUCTURES AND BRAINSTEM - p. 329

Dzeman M.I.
MONOGRAPH "FEDIR STEPANOVYCH
TSYTSURIN: A VIEW THROUGH
THE CENTURIES (DISCOURSE ABOUT THE
SUCCESSION OF CLINICAL MEDICINE TRADITIONS
DEDICATED TO THE 200TH ANNIVERSARY
OF THE KYIV INTERNISTS SCHOOL FOUNDER) - p. 330

Мельник В. М.
ПЕРВАЯ УКРАИНСКАЯ
КНИГА О СОЦИОКУЛЬТУРНОЙ
АНТРОПОЛОГИИ - p. 332

Пивоваров А.А.
ЗАНИМАТЕЛЬНАЯ АСТРОНОМИЯ
В НОВЕЛЛАХ ТЕСОВОГО
ХАРАКТЕРА: УЧЕБНОЕ ПОСОБИЕ – p. 334

Shikhalieva S.
SPECIFICS OF THE PARONYMY
AND METHODS OF ITS TEACHING – p. 335

Kazarezov Mikhail
«RECONSTRUCTIVE SURGERY
OF THE COMPLICATED DAMAGES» – p. 336

Кривицкий Б.Х.
«УЧЕБНЫЕ ЭЛЕКТРОННЫЕ
СРЕДСТВА В ВУЗЕ» – p. 337

VIENNA, AUSTRIA

February, 2016

THE EUROPEAN ASSOCIATION OF PEDAGOGUES AND PSYCHOLOGISTS "SCIENCE"



The European Association of pedagogues and psychologists

“Science”

International scientific periodical journal

“THE UNITY OF SCIENCE”

Vienna, Austria, 2016

Single photocopies of single chapters may be made for personal use as allowed by national copyright laws. Permission of the Publisher and payment of a fee is required for all other photocopying, including multiple or systematic copying, copying for advertising or promotional purposes, resale, and all forms of document delivery. Special conditions are available for educational institutions that wish to make photocopies for non-profit educational classroom use. Permission of the Publisher is required for all other derivative works, including compilations and translations. Electronic Storage or Usage Permission of the Publisher is required to store or use electronically any material contained in this work, including any chapter or part of a chapter. Except as outlined above, no part of this work may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without prior written permission of the Publisher.

International scientific professional periodical journal "THE UNITY OF SCIENCE" / publishing office Friedrichstrabe 10– Czech Republic – Prague, 2015

"The Unity of science" is the international scientific professional periodical journal which includes the scientific articles of students, graduate students, postdoctoral students, doctoral candidates, research scientists of higher education institutions of Europe, Russia, the countries of FSU and beyond, reflecting the processes and the changes occurring in the structure of present knowledge. It is destined for teachers, graduate students, students and people who are interested in contemporary science.

The editorial board

Alexander Pacsr
Astakhov Vladimir
Batsyleva Olga
Buriak Alexander
Emma Kahl
George Berins
Georgieva Sonia
Hans Jensen
Helen Zag
Homyuk Victor
Ilyina Ekaterina
Ivanova Svetlana
Ivanova Zoya
Jakob Haberzettl
Kaydalova Lydia
Kovtun Elena
Logvina Svetlana
Lokhvitskaya Lyubov

Lucius Calle
Lukas Adlerflycht
Lyakisheva Anna
Marek Szymański
Mariusz Bashinsky
Matra Lentz
Matthias Haenlein
Mukanova Sulesh
Patrick Schuppe
Petkova Ekaterina
Petruk Vera
Prokofieva Olesya
Rymarenko Nataliya
Sergienko Konstantin
Sophia Lai
Stanislaw Krauchensky
Stetsenko Iryna
Vaskes Abanto Hesus Estuardo
Yakobchuk Elena

Designed by Mr. D. Heisenberg

Publishing office Beranových str., 130 of.37, Prague, Czech Republic, 2015
[www. eapps.info](http://www.eapps.info)



© 2016 The European Association of pedagogues and psychologists Science”, Vienna, Austria

© 2016 Article writers

© 2016 All rights reserved

ASSOCIATION OF MARKERS OF THE SYSTEM OF FIBRINOLYSIS OF THE BLOOD PLASMA WITH A/C POLYMORPHISM OF DIO1 AND PRO197LEU OF GPX1 GENES IN PATIENTS WITH NONVIRAL CHRONIC HEPATITIS AND LIVER CIRRHOSIS.....276

Chympoy K.A.
Higher State Educational Establishment of Ukraine
«Bukovinian State Medical University»,
Assistant, Department of Internal Medicine

Pavlyukovich N.D.
Higher State Educational Establishment
of Ukraine «Bukovinian State Medical University»,
Assistant, Department of Internal Medicine,
clinical pharmacology and occupational diseases

Buriak O.G.
Higher State Educational Establishment
of Ukraine «Bukovinian State Medical University»,
Associate Professor, Department of Pediatrics,
neonatology and perinatal medicine

DYNAMICS OF CHANGES IN THE DEGREE OF DEPOLARIZATION OF LASER RADIATION, MULTIPLE BY THE MYOCARDIAL TISSUE, FOR DETERMINATION OF THE DEATH TIME.....279

Bachynskyy Viktor
Department of Forensic Medicine and Medical Law
Higher State Educational Establishment of Ukraine
“Bukovinian State Medical University”

Pavlyukovich Natalia
Department of Internal Medicine,
Clinical Pharmacology and Occupational Diseases
Higher State Educational Establishment of Ukraine
“Bukovinian State Medical University”

Pavlyukovich Alexandr
Department of Forensic Medicine and Medical Law
Higher State Educational Establishment of Ukraine
“Bukovinian State Medical University”

ERYTHROCYTE AT CONCOMITANT COURSE OF CORONARY ARTERY DISEASE AND ANEMIA: PHYSICAL VIEW ON CLINICAL ASPECTS OF COMBINED PATHOLOGY.....283

Natalia Pavlyukovich
Department of Internal Medicine,
Clinical Pharmacology and Occupational Diseases Higher
State Educational Establishment of Ukraine “Bukovinian State
Medical University”

Kristina Chimpoi
Department of Internal Medicine
Higher State Educational Establishment of Ukraine
“Bukovinian State Medical University”

Pavlyukovich Alexandr
Department of Forensic Medicine
and Medical Law Higher State Educational Establishment
of Ukraine “Bukovinian State Medical University”

Kozar Marta
Cardiological Department
Chernivtsi Regional Hospital for Disabled
Soldiers of the Patriotic War

THE CASE OF SYNDROME LEJEUNE IN CHILD.....288



Ryznychuk M.O.
Assistant Department of Pediatrics
and Medical Genetics, Ph.D.
Higher educational institution of Ukraine
“Bukovina State Medical University”



Polishchuk M.I.
PhD, head of the of organization
and development of treatment and preventive
care of the Department of Health
Chernivtsi Regional State Administration



Lastivka I.V.
Associate Professor of Pediatrics
and Medical Genetics, Ph.D.
Higher educational institution Ukraine
“Bukovina State Medical University”

ASSOCIATION OF MARKERS OF THE SYSTEM OF FIBRINOLYSIS OF THE BLOOD PLASMA WITH A/C POLYMORPHISM OF DIO1 AND PRO197LEU OF GPX1 GENES IN PATIENTS WITH NONVIRAL CHRONIC HEPATITIS AND LIVER CIRRHOSIS

КОРЕКЦІЯ МЕКСИДОЛОМ ФУНКЦІОНАЛЬНОГО СТАНУ НИРОК ЗА УМОВ РОЗВИТКУ ПЕРШОЇ СТАДІЇ ГАРЯЧКИ

Chympoy K.A.

Higher State Educational Establishment of Ukraine
«Bukovinian State Medical University»,
Assistant, Department of Internal Medicine
E-mail: chimpoik@mail.ru

Buriak O.G.

Higher State Educational Establishment of Ukraine
«Bukovinian State Medical University»,
Associate Professor, Department of Pediatrics,
neonatology and perinatal medicine
E-mail: buriak@neonatology.in.ua

Pavlyukovich N.D.

Higher State Educational Establishment of Ukraine
«Bukovinian State Medical University»,
Assistant, Department of Internal Medicine,
clinical pharmacology and occupational diseases
E-mail: natasha.pavlyukovich@gmail.com

Abstract. To study peculiarities of the indices of the fibrinolysis in patients with chronic diffuse liver diseases depending A/C polymorphism in DIO1 and GPX1 Pro197Leu gene. Estimation of the indices of fibrinolytic blood activity in patients with chronic diffuse liver disease showed that total fibrinolytic activity and enzymatic fibrinolytic activity were reliably lower than that of the control indices at the same time non-enzymatic fibrinolytic activity increased and did not reveal their dependence on the distribution of A/C polymorphism of DIO1 gene and Pro197Leu polymorphism of GPX1.

Key words: chronic diffuse liver disease, polymorphism, fibrinolysis.

Introduction. The analysis of genetic associations plays an important role in the examination of the role of genetic factors involved in the development of polymorphic diseases, and chronic diffuse liver disease in particular [2, 4, 5, 9]. The difference of marker allele frequency in patients with certain pathology and healthy individuals gives the evidence to draw a conclusion about the link between a particular allele and corresponding pathology [3, 6]. The information available concerning the links of chronic diffuse liver disease pathogenesis allows detecting the range of genes-candidates which potential relation with this pathology needs further investigation [6, 8].

Due to recent scientific research both of Ukrainian and foreign scientists the concept of relations between indices of fibrinolysis and expression of various genes is beyond any doubt [1,7,8]. Although dependence of the above indices upon A/C polymorphism DIO1 and Pro197Leu GPX1 gene in patients with chronic diffuse liver disease remains above the attention of researchers.

The aim the study. To study peculiarities of the indices of the fibrinolysis in patients with chronic diffuse liver diseases depending on A/C polymorphism in DIO1 and GPX1 Pro197Leu gene.

Materials and methods. 28 patients with chronic diffuse liver disease aged from 34 to 72 were examined. Depending on the distribution of DIO1 gene A/C polymorphism the patients were divided into three groups: AA-genotype carriers – 9 patients, AC-genotype – 11, CC-gen-

otype – 8. Depending on GPX1 gene Pro197Leu polymorphism there were 12 homozygotes by Pro-allele, 8 – by Leu-allele and 8 ProLeu-heterozygotes.

The diagnosis of chronic diffuse liver disease was made on the basis of anamnesis, generally accepted complex of clinical-laboratory and instrumental investigation methods, USD of the abdominal organs. Patients with chronic hepatitis and cirrhosis of a viral etiology, Wilson-Konovalov disease, congenital insufficiency of α -antitripsin (α -inhibitor of proteinase), idiopathic (genetic) hemochromatosis, autoimmune hepatitis were excluded from the study.

Alleles of A/C regions in DIO1 gene and Pro197Leu in GPX1 gene were studied by means of excretion of genome DNA from leukocytes of the peripheral blood with further amplification of a polymorphic region by means of polymerase chain reaction (PCR) on the programmed amplification «Amply-4L» («Biocom», Moscow) with individual temperature program for the parameters of every gene. Table 1 presents succession of oligonucleotides in primers and their calculation positions on chromosomes.

DNA extraction was conducted by means of «DNA-sorb-B» reagents, variant 100 (Russian) according to the instruction. Purified DNA was kept under the temperature of $20 \pm 2^\circ\text{C}$. Samples for PCR were prepared by means of «AmplySense – 200 – 1» set (Russian). Bcl I restriction endonuclease produced by «SibEnzyme» firm (Russian) was used to discriminate DIO1 gene alleles.

Table 1

Succession of oligonucleotides in primers used for polymerase chain reaction (PCR) to identify A/C polymorphism of DIO1 gene and Pro197Leu of GPX1 gene

DIO1	1 p33-p32	Direct	5'-GAACTTGATGTGAAGGCTGGA-3'
		Reverse	5'-TAACCTCAGCTGGGAGTTGTTT-3'
GPX1	3p21	Direct	5'-TCGAAGCCCTGCTGTCTCA-3'
		Reverse	5'-CGAGACAGCAGCACTGCAA-3'

Total non-enzymatic and enzymatic fibrinolysis of citrated blood plasma was estimated by asofibrinolysis (Simko Ltd., Ukraine).

The results obtained are calculated by means of Biostat program with the use of Student t-criterion.

Results and discussion. The indices of fibrinolysis in patients with chronic diffuse liver disease did not experience reliable changes depending on polymorphism of DIO1 gene and were statistically different from the group of practically healthy individuals (table 2).

Examination of fibrinolytic blood activity showed that total fibrinolytic activity of the blood plasma in all the group of patients was reliably lower than that of the control indices: CC-genotype – on 19% (P1<0,001), AC-genotype and AA-genotype – on 22,1% and 18,4% correspondingly (P1<0,001) without reliable intergroup difference. At the same time non-enzymatic fibrinolytic activity in all groups of patients increases in comparison with the control on 37,3% (P1<0,001), 33,3% and 31,4% correspondingly (P1<0,001). Total fibrinolytic activity index in patients with AA-genotype was reliably lower than that of the control in 1,7 times (P1<0,001), CC-genotype – in 1,8 times (P1<0,001), while for the patients with AC-genotype a maximal inhibition of total fibrinolytic activity was registered – in 1,9 times (P1<0,001).

lower than that of the control values: in patients with ProPro-genotype – on 22,1% (P<0,001), with ProLeu-genotype and LeuLeu-genotype – on 19,6% (P<0,001) and 18,4% (P<0,01) respectively without reliable difference between the groups.

Non-enzymatic fibrinolytic activity in patients of all the groups elevated, and increasing of this index in comparison with the control group was indicative of it: on 25,5% (P<0,001), 35,3% (P<0,001), and 41,2% (P<0,001) in the carriers of ProPro-, ProLeu- and LeuLeu-genotype respectively.

Thus, A/C polymorphism of DIO1 gene and Pro197Leu polymorphism of GPX1 gene does not influence upon the indices of the system of fibrinolysis in patients with chronic diffuse liver diseases.

Received results can be explained by the increasing of the free radical oxidation processes due to the liver-cell failure. Free radicals are able to directly destroy NO [6]. Increased lipid peroxidation of membranes is the result of damaging the structure of the endothelium and its violation NO-producing ability. Developing an absolute or relative deficiency of NO, is required for normal regulation of vascular tone. The weakening of the NO-dependent vasodilatory reactions, in its turn, leads to increased vascular tone, increased blood clots, and as a result, tissue hypoxia. In addition, the reduced inhibitory effect of NO on plate-

Table 2

Indices of the fibrinolysis in patients with chronic diffuse liver disease depending on A/C polymorphism of DIO1 gene (M±m)

Total fibrinolytic activity, mcmolazofibrin/1mL per hour	1,63±0,041	1,33±0,072 P ₁ <0,001	1,27±0,047 P ₁ <0,001 P ₂ >0,05	1,32±0,050 P1<0,001 P2>0,05 P3>0,05
Non-enzymatic fibrinolytic activity, mcmolazofibrin/1mL per hour	0,51±0,019	0,67±0,038 P ₁ <0,001	0,68±0,029 P ₁ <0,001 P ₂ >0,05	0,70±0,020 P1<0,001 P2>0,05 P3>0,05
Enzymatic fibrinolytic activity, mcmolazofibrin/1mL per hour	1,12±0,051	0,66±0,084 P ₁ <0,001	0,59±0,072 P1<0,001 P2>0,05	0,62±0,065 P1<0,001 P2>0,05 P3>0,05

Notes: n – numbers of observations; P1 – probability of changes concerning the control; P2 – probability of changes concerning the group of patients with AA-genotype; P3 – probability of changes concerning the group of patients with AC-genotype

Table 3 presents the results of examination of fibrinolysis in patients with chronic diffuse liver disease depending on the distribution of Pro-197Leu polymorphism of GPX1 gene.

Examination of fibrinolytic blood activity showed that total fibrinolytic activity of the blood plasma in patients of all the groups was reliably

let aggregation, leukocyte adhesion to the endothelium and smooth muscle cell proliferation of the vascular wall, creates prerequisites for vascular disorders.

Endothelial cells are involved in various physiological processes, such as vascular tonus regulation, hemostasis, fibrinolysis, inflamma-

Table 3

Indices of the fibrinolysis in patients with chronic diffuse liver diseases depending on Pro197Leu polymorphism of GPX1 gene (M±m)

Total fibrinolytic activity, mcmolazofibrin/1mL per hour	1,63±0,041	1,27±0,049 P ₁ <0,001	1,31±0,062 P ₁ <0,001 P ₂ >0,05	1,33±0,055 P ₁ <0,01 P ₂ >0,05 P ₃ >0,05
Non-enzymatic fibrinolytic activity, mcmolazofibrin/1mL per hour	0,51±0,019	0,64±0,034 P ₁ <0,001	0,69±0,022 P ₁ <0,001 P ₂ >0,05	0,72±0,024 P ₁ <0,001 P ₂ >0,05 P ₃ >0,05
Enzymatic fibrinolytic activity, mcmolazofibrin/1mL per hour	1,12±0,051	0,61±0,083 P ₁ <0,001	0,68±0,077 P ₁ <0,001 P ₂ >0,05	0,56±0,074 P ₁ <0,001 P ₂ >0,05 P ₃ >0,05

Notes: n – numbers of observations;

P1 – probability of changes concerning the control

P2 – probability of changes concerning the group of patients with ProPro-genotype

P3 – probability of changes concerning the group of patients with ProLeu-genotype

tion. They maintain the barrier permeability for exchange and active transport of substances in the arterial wall. Endothelial dysfunction, caused by the pathological mechanisms, including oxidative stress and increased cell adhesion, obviously inhibits the fibrinolytic activity of blood in the examined patients. Particularly, statistically significant decreasing of total fibrinolytic activity for 20.2% due to decreasing of the fibrin enzyme-related lysis for 45,5% along with the increasing of non-productive non-enzymatic fibrinolytic activity for 35, 2% were detected in the patients with chronic diffuse diseases of the liver.

Conclusions:

1. Examination of the indices of fibrinolytic blood activity in patients

with nonviral chronic hepatitis and liver cirrhosis showed that total fibrinolytic activity for 20,2% and enzymatic fibrinolytic activity was reliably lower for 45,5% than that of the control indices at the same time non-enzymatic fibrinolytic activity increases for 35,2%.

2. Pro197Leu polymorphism of GPX1 gene and A/C polymorphism of DIO1 gene does not influence upon the indices of the system of fibrinolysis in patients with nonviral chronic hepatitis and liver cirrhosis.

The prospects of proceeding investigations will be further studies of pathogenetic peculiarities chronic diffuse liver diseases with the aim to find the mechanisms of their occurrence and progress and substantiation of the improved methods to correct and prevent the given pathology.

References

1. Bastaki M. Genotype-activity relationship for Mn-superoxide dismutase, glutathione peroxidase-1 and catalase in humans / M. Bastaki, K. Huen, P. Manzanillo [et al.] // *Pharmacogenetics and Genomics*. – 2006. – Vol. 16. – P. 279-286.
2. Chrissobolis S. Glutathione peroxidase-1 plays a major role in protecting against angiotensin II-induced vascular dysfunction / S. Chrissobolis, S.P. Didion, D.A. Kinzenbaw [et al.] // *Hypertension*. – 2008. – Vol. 51. – P. 872-877.
3. Chympoy K.A. Kharakterystyka tytriv antytil do tkany shchytovidnoyi zalozy, pokaznykiv tyreoidnoho homeostazu, pro- ta antyoksydantnoyi system krovu u khvorykh na khronichni dyfuzni zakhvoryuvannya pechinky zalezho vid polimorfizmu heniv / K.A. Chympoy, N.V. Pashkovska, A.I. Kurchenko // *Imunohiyya ta alerholiyya*. – 2010. – № 3-4. – P. 131-136.
4. Eremina E.YU. Faktory riska progressirovaniya hronicheskikh gepatitov i cirrozov pecheni / E.YU. Eremina // *EHKsperimental'naya i klinicheskaya gastroenterologiya*. – 2008. – № 6. – P. 101-106.
5. Hodge A. Coagulopathy in liver disease: The whole is greater than the sum of its parts / A. Hodge, P. Crispin // *J. of Gastroenterol. and Hepatol.* – 2010. – Vol. 25, № 1. – P. 1-2.
6. Ivashkin V.T. Klinicheskaya gepatologiya segodnya i zavtra / V.T. Ivashkin, A.O. Bueverov // *Ros. zhurn. gastroenterologii, gepatologii, koloproktologii*. – 2005. – T. 12., № 1. – P. 4-9.
7. Pavlov CH.S. Sovremennye predstavleniya o patogeneze, diagnostike i lechenii fibroza pecheni / CH.S. Pavlov, YU.O. SHul'pekova, V.B. Zolotarevskij, V.T. Ivashkin // *Ros. zhurn. gastroenterol., gepatol., koloproktol.* – 2005. – T. 15, № 2. – P. 13-20.
8. Sneddon A.A. Cytokine-induced monocyte adhesion to endothelial cells involves platelet-activating factor: Suppression by conjugated linoleic acid / A.A. Sneddon, E. McLeod, K.W. Wahle, J.R. Arthur // *Biochimica et Biophysica Acta-Molecular and Cell Biology of Lipids*. – 2006. – Vol. 761. – P. 793-801.
9. Wagner Andreas H. Upregulation of Glutathione Peroxidase Offsets Stretch-Induced Proatherogenic Gene Expression in Human Endothelial Cells Arteriosclerosis, Thrombosis, and Vascular Biology / H. Wagner Andreas, O. Kautz, K. Fricke [et al.] // *ATVBAHA*. – 2009. – Vol. 21. – P. 316-325.