

diabetes and careful monitoring of blood glucose levels, especially in the acute period of stroke, to prevent an increase in the area of the lesion and recurrent stroke.

Dobriansky V. V.

RISK FACTORS FOR RECURRENT HYPOGLYCEMIA IN PATIENTS WITH TYPE II DIABETES MELLITUS IN THE PREHOSPITAL STAGE

Department of Anesthesiology and Intensive Care

Bukovynian State Medical University

About 500 million people worldwide have diabetes, about 90% of whom have type II diabetes. According to WHO statistics, in 2019, more than 1.5 million deaths were caused by diabetes and its complications, among which one of the most common is hypoglycemia. In most patients receiving insulin, hypoglycemia develops with varying frequency; in 30% of cases, there is severe hypoglycemia. According to the Center of Emergency Medical Care and Disaster Medicine (CEMCDM) in the Chernivtsi region, the number of calls about hypoglycemia is 10-15% of the total number of calls per year. A lot of them are complicated and recurrent, so it is important to understand and study the risk factors of recurrent hypoglycemia in a prehospital setting.

Purpose: to investigate and analyze the risk factors for recurrent hypoglycemia in patients with type II diabetes. 33 ambulance call cards and the annual report on the work of CEMCDM of Chernivtsi were analyzed. The results of general clinical and laboratory research methods are studied, the general methods of statistics are used.

The sample consisted of 33 patients with type II diabetes who were on insulin therapy, of which 19 (57.5%) were men and 14 (42.5%) were women aged from 17 to 76 years. The moderate diabetes was detected in 16 people (48.5%), mild - in 10 (30.3%), and severe - in 7 patients (21.2%). Among those surveyed, the glycemic level was on average 2.27 mmol/l (0.5-2.9 mmol/l). Aftercare, the average blood glucose level was 7.3 mmol/L. There were 3 cases (9%) of very severe hypoglycemia (less than 1.0 mmol/l). This group of patients was characterized by risk factors such as old age, prolonged diabetes, cardiovascular disease, difficulty recognizing symptoms and alcohol abuse, glucose resistance, and recurrent episodes in anamnesis. Repeated calls accounted for 21.2% of the total number of call cards analyzed, most of them for patients with severe and moderate diabetes who were on insulin therapy. For these patients, in addition to standard treatment (administration of 80-100 ml of 40% glucose), 200 ml of 5% glucose was added. The average time of arrival of an ambulance (TAA) was 28 minutes, including 17.4 minutes in the city and 41.1 minutes in the countryside. A direct correlation was found between TAA and the amount of 40% glucose administered ($r = 0.6$). It was also found that 66.6% of respondents abused alcohol. The incidence of severe hypoglycemia was higher in patients with severe diabetes and who did not follow a diet. In women, the incidence was slightly lower than in men, but they had lower glucose levels compared with men ($p < 0.05$).

The risk of recurrent hypoglycemia is high in patients with older age, long and severe diabetes, comorbidities, in people who abuse alcohol. A direct correlation was established between the NPV and the amount of 40% glucose administered. Thus, it is important to provide timely care for hypoglycemia in patients with type II diabetes, as well as hospitalization of patients with the above risk factors to specialized hospitals.

Malaiko S.S.

FREQUENCY OF PULMONARY EDEMA DEVELOPMENT IN PATIENTS WITH ACUTE CORONARY SYNDROME AND DIABETES TYPE 2

Department of Anesthesiology and Intensive Care

Bukovynian State Medical University

Pulmonary edema is a liquid accumulation in the tissue and air spaces of the lungs. It leads to impaired gas exchange and may cause respiratory failure. It is due to either failure of the left ventricle of the heart to remove blood adequately from the pulmonary circulation. On the basis of the data of the register of acute coronary syndrome to study the incidence of pulmonary edema that