

**МІНІСТЕРСТВО ОХОРОНИ ЗДОРОВ'Я УКРАЇНИ  
БУКОВИНСЬКИЙ ДЕРЖАВНИЙ МЕДИЧНИЙ УНІВЕРСИТЕТ»**



## **МАТЕРІАЛИ**

**105-ї підсумкової науково-практичної конференції  
з міжнародною участю  
професорсько-викладацького персоналу  
БУКОВИНСЬКОГО ДЕРЖАВНОГО МЕДИЧНОГО УНІВЕРСИТЕТУ  
присвяченої 80-річчю БДМУ  
05, 07, 12 лютого 2024 року**

Конференція внесена до Реєстру заходів безперервного професійного розвитку,  
які проводитимуться у 2024 році № 3700679

**Чернівці – 2024**

УДК 001:378.12(477.85)

ББК 72:74.58

М 34

Матеріали підсумкової 105-ї науково-практичної конференції з міжнародною участю професорсько-викладацького персоналу Буковинського державного медичного університету, присвяченої 80-річчю БДМУ (м. Чернівці, 05, 07, 12 лютого 2024 р.) – Чернівці: Медуніверситет, 2024. – 477 с. іл.

ББК 72:74.58

У збірнику представлені матеріали 105-ї підсумкової науково-практичної конференції з міжнародною участю професорсько-викладацького персоналу Буковинського державного медичного університету, присвяченої 80-річчю БДМУ (м. Чернівці, 05, 07, 12 лютого 2024 р.) із стилістикою та орфографією у авторській редакції. Публікації присвячені актуальним проблемам фундаментальної, теоретичної та клінічної медицини.

Загальна редакція: професор Геруш І.В., професорка Грицюк М.І., професор Безрук В.В.

Наукові рецензенти:

професор Братенко М.К.

професор Булик Р.Є.

професор Гринчук Ф.В.

професор Давиденко І.С.

професор Дейнека С.Є.

професорка Денисенко О.І.

професор Заморський І.І.

професорка Колоскова О.К.

професор Коновчук В.М.

професор Пенішкевич Я.І.

професорка Хухліна О.С.

професор Слободян О.М.

професорка Ткачук С.С.

професорка Годоріко Л.Д.

професор Юзько О.М.

професорка Годованець О.І.

ISBN 978-617-519-077-7

© Буковинський державний медичний  
університет, 2024

**Kushnir O.Yu.**

**THE STATE OF THE HEXOSOMONOPHOSPHATE SHUNT IN THE LIVER OF RATS WITH LATENT ALLOXAN DIABETES UNDER CONDITIONS OF MELATONIN ADMINISTRATION AND ALTERED PHOTOPERIOD**

*Department of Bioorganic and Biological Chemistry and Clinical Biochemistry  
Bukovinian State Medical University*

**Introduction.** The pentose phosphate pathway (or hexosomonophosphate shunt) is necessary for transformed cells to synthesize ribonucleotides and is the main source of NADPH<sub>2</sub>.

**The aim of the study.** To find out the influence of melatonin on the activities of glucose-6-phosphate dehydrogenase (G-6-PDH), 6-phosphogluconate dehydrogenase (6-PGDH) and transketolase (TK) in the liver of rats with latent diabetes under conditions of normal, hyper- and hypofunction of the pineal gland.

**Material and methods.** Experiments were conducted on male outbred white rats weighing 180±10 mg. Photoperiodic changes were simulated for 12 days: 1) natural equinox from March 19 to 25 (L12:D12); 2) constant light during the day (500 lux); 3) constant darkness during the day. Alloxan diabetes was induced by intraperitoneal administration of a 5% solution of alloxan monohydrate at a dose of 170 mg/kg once, after a 24-hour fast. Basal glycemia (BG) was determined using the One Touch Ultra device (Johnson & Johnson, USA). On the third day, the death of 50% of diabetic rats was observed. Experimental animals were divided into the following groups: 1) intact rats (control); 2) rats with latent diabetes (BG≤6.9 mmol/l); 3) rats with latent diabetes mellitus, which were intraperitoneally injected with melatonin (Sigma, USA) at a dose of 10 mg/kg of body weight from the 5th day after the administration of alloxan for a week at 8:00 a.m. daily. Animals were euthanized on the 12th day from the beginning of the experiment in accordance with the ethical principles of experiments on animals. In the supernatant obtained after centrifugation of 5% liver homogenate at 900 g, enzyme activities were determined according to standard methods. Statistical processing was performed according to the Student.

**Results.** The activities of G-6-PDH, 6-PGDH in animals with latent diabetes were 77 and 64% higher, respectively, than in control. In the liver of control animals that were under conditions of constant darkness, the activities of G-6-PDG, 6-PGDH, and TK increased by 38, 29, and 12%, respectively, under conditions of constant light, the activities of G-6-PDG, 6-PGDG, and TK decreased by 18, 10 and 15%, respectively, in comparison with the indicators of intact animals under equinox conditions. With constant darkness against the background of latent diabetes, the activities of G-6-PDG, 6-PGDH and TK increased by 115, 95 and 30%, respectively; under conditions of constant light, the activities of G-6-PDG, 6-PGDH, and TK were 24, 16, and 22% lower, respectively, than in the group of intact animals under equinox conditions. But if oxidative stress develops in the cell, then the pentose phosphate pathway will "reciprocate" with glycolysis through fructose-6-phosphate, which is formed in the process of the non-oxidative phase. If the cell is in a state of oxidative stress, then it immediately stops glycolysis and transfers all glucose to the pentose phosphate pathway the path for the synthesis of NADPH. Administration of melatonin to animals with latent diabetes under all lighting conditions led to the normalization of all investigated parameters (except for the activity of G-6-PDH under conditions of constant light in latent diabetes, which increased by 20% compared to intact animals at the equinox).

**Conclusions.** Melatonin allows to restore the violation of carbohydrate metabolism in the context of the restoration of biosynthetic processes and antioxidant protection due to the regulation of the enzyme activity of the hexosomonophosphate shunt of glucose-6-phosphate oxidation.

**Lenha E.L.**

**EFFECT OF MELATONIN ON CATALASE ACTIVITY IN THE BLOOD OF RATS**

*Department of Bioorganic and Biological Chemistry and Clinical Biochemistry  
Bukovinian State Medical University*

**Introduction.** Violation of the body's antioxidant resistance under the influence of physical, chemical or stressful factors requires early diagnosis with subsequent correction to prevent and