



p-ISSN 2224-0721
e-ISSN 2307-1427

International journal of endocrinology

МІЖНАРОДНИЙ ЕНДОКРИНОЛОГІЧНИЙ ЖУРНАЛ

SANOFI DIABETES


Том 13, № 4, 2017

СПЕЦІАЛІЗОВАНІЙ РЕЦЕНЗОВАНИЙ
НАУКОВО-ПРАКТИЧНИЙ ЖУРНАЛ



ЗАСЛАВСКИЙ
Издательский дом

Буковинський державний медичний університет



Міжнародний ендокринологічний журнал

Международный эндокринологический журнал

International journal of endocrinology

Спеціалізований рецензований науково-практичний журнал
Заснований у вересні 2005 року
Періодичність виходу: 8 разів на рік

Том 13, № 4, 2017

Включений в наукометричні та спеціалізовані бази даних Ulrichsweb Global Serials Directory, Index Copernicus, Directory of Research Journals Indexing (DRJI), WorldCat, РІНЦ (Science Index), Google Scholar, «Джерело», «Наукова періодика України», Academic Resource Index (Research Bible), «КіберЛенінка», НБУ ім. В.І. Вернадського, CrossRef, Universal Impact Factor, General Impact Factor, International Committee of Medical Journal Editors (ICMJE), SHERPA/RoMEO, Bielefeld Academic Search Engine (BASE)

Імпакт-фактор РІНЦ: 0,227

Index Copernicus ICV: 61.19



Міжнародний ендокринологічний журнал

Спеціалізований рецензований
науково-практичний журнал

Том 13, № 4, 2017

DOI: 10.22141/2224-0721.13.4.2017
p-ISSN 2224-0721
e-ISSN 2307-1427

Передплатний індекс: 94553



Співзасновники:

Буковинський державний медичний університет,
Заславський О.Ю.

Завідуюча редакцією
Купріненко Н.В.

Адреса для звертань:

Із питань передплати:

info@mif-ua.com,
тел. +38 (044) 223-27-42,
+38 (067) 325-10-26

З питань розміщення реклами
та інформації про лікарські засоби:

reclama@mif-ua.com,
office@zaslavsky.kiev.ua
golubnichayan@gmail.com

Журнал внесено до переліку наукових фахових видань України, в яких можуть публікуватися результати дисертаційних робіт на здобуття наукових ступенів доктора і кандидата наук. Наказ МОН України від 15.04.2014 р. № 455.

Рекомендується до друку та до поширення через мережу Інтернет вченого радою Вищого державного навчального закладу ІV рівня акредитації «Буковинський державний медичний університет» МОЗ України (22 червня 2017 р., протокол № 12).

Українською, російською та англійською мовами

Свідоцтво про державну реєстрацію друкованого засобу масової інформації КВ № 19313-9113ПР. Видано Державною реєстраційною службою України 06.09.2012 р.

Формат: 60×84/8. Ум. друк. арк. 12,32
Зам. 2017-іє-84. Тираж 3 000 прим.

Адреса редакції:
а/с 74, м. Київ, 04107, Україна
Тел./факс: +38 (044) 223-27-42
E-mail: medredactor@i.ua

(Тема: До редакції «Міжнародного
ендокринологічного журналу»)
www.mif-ua.com, http://iej.zaslavsky.com.ua

Видавець Заславський О.Ю.
Адреса для листування: а/с 74, м. Київ, 04107,
Адреса реєстрації: пр. Ленінський, 25/126,
м. Донецьк, 83102

Свідоцтво суб'єкта видавничої справи
ДК № 2182 від 13.05.2005 р.

Друк: ООО «Ландпрес»
ул. Алчевських, 2, г. Харків, 61002

Головний редактор
Володимир Іванович ПАНЬКІВ

Науковий редактор
Тарас Миколайович БОЙЧУК

Редакційна колегія

- | | |
|--------------------------------|----------------------------------|
| Авраменко Т.В. (Київ) | Маньківський Б.М. (Київ) |
| Бобирьова Л.Є. (Полтава) | Мітченко О.І. (Київ) |
| Боднар П.М. (Київ) | Могілевський С.Ю. (Київ) |
| Большова О.В. (Київ) | Пасечко Н.В. (Тернопіль) |
| Бондаренко В.О. (Харків) | Поворознюк В.В. (Київ) |
| Вернигородський В.С. (Вінниця) | Пашковська Н.В. (Чернівці) |
| Веселовська З.Ф. (Київ) | Перцева Т.О. (Дніпро) |
| Власенко М.В. (Вінниця) | Полторак В.В. (Харків) |
| Генделека Г.Ф. (Одеса) | Резников О.Г. (Київ) |
| Гончарова О.А. (Харків) | Сергієнко О.О. (Львів) |
| Зелінська Н.Б. (Київ) | Сидорчук Л.П. (Чернівці) |
| Іващук О.І. (Чернівці) | Сіренко Ю.М. (Київ) |
| Караченцев Ю.І. (Харків) | Скрипник Н.В. (Івано-Франківськ) |
| Кирилюк М.Л. (Київ) | Соколова Л.К. (Київ) |
| Козаков О.В. (Харків) | Томашевський Я.І. (Львів) |
| Комісаренко Ю.І. (Київ) | Тронько М.Д. (Київ) |
| Корпачов В.В. (Київ) | Хижняк О.О. (Харків) |
| Кравченко В.І. (Київ) | Черенько С.М. (Київ) |
| Кравчун Н.О. (Харків) | Юзвенко Т.Ю. (Київ) |
| Ларін О.С. (Київ) | |
| Лучицький Є.В. (Київ) | |

Редакційна рада

- | | |
|------------------|------------------------|
| Аметов О.С. | Шестакова М.В. |
| | (Москва, Росія) |
| Арістархов В.Г. | Dr. Agaç F. |
| | (Рязань, Росія) |
| Базарбекова Р.Б. | Alekna V. |
| | (Алмати, Казахстан) |
| Валєєва Ф.В. | Czupryniak L. |
| | (Варшава, Польща) |
| Вербовой А.Ф. | Ferrannini E. |
| | (Піза, Італія) |
| Данилова Л.І. | Holick M.F. |
| | (Бостон, США) |
| Дедов І.І. | Mascarenhas M.R. |
| | (Лісабон, Португалія) |
| Зельцер М.Ю. | Mota M. |
| | (Крайова, Румунія) |
| Ісмаїлов С.І. | Nikberg I. |
| | (Мельбурн, Австралія) |
| Мамедов М.Н. | Radzeviciene L. |
| | (Каунас, Литва) |
| Мельниченко Г.А. | Rurik Imre |
| | (Дебрецен, Угорщина) |
| Мірзазаде В. | Standl E. |
| | (Мюнхен, Німеччина) |
| Мохорт Т.В. | Szabolcs I. |
| | (Будапешт, Угорщина) |
| Романчишен А.П. | Taton J. |
| | (Варшава, Польща) |
| Свириденко Н.Ю. | Ткач Іван |
| | (Кошице, Словаччина) |
| Трошина К.А. | Yki-Järvinen H. |
| | (Гельсінкі, Фінляндія) |
| | Zgliczynski S. |
| | (Варшава, Польща) |

Відповідальний секретар Іван Іваносич ПАВЛУНИК

Редакція не завжди поділяє думку автора публікації. Відповідальність за вірогідність фактів, власних імен та іншої інформації, використаної в публікації, несе автор. Передрук та інше відтворення в якій-небудь формі в цілому або частково статей, ілюстрацій або інших матеріалів дозволені тільки при попередній письмовій згоді редакції та з обов'язковим посиланням на джерело. Усі права захищені.

© Буковинський державний медичний університет, 2017
© Заславський О.Ю., 2017

Практикуючому ендокринологу

- Що нового у виявленні та лікуванні дисліпідемії і профілактиці прогресування атеросклерозу? 35

Огляд літератури

*Сорокман Т.В., Молдован П.М.,
Попелюк Н.О., Макарова О.В.*

- Генетична основа ендокринної патології 47

*Оленович О.А., Пашковська Н.В.,
Пашковський В.М., Бойко І.В.*

- Ураження периферичної нервової системи при гіпотиреозі: сучасний погляд на проблему (огляд літератури) 57

Клінічна тиреоїдологія

*Вернигородский В.С., Фетисова Н.М.,
Вернигородская М.В.*

- Патогенетическое значение провоспалительных цитокинов и дислипидемии в развитии сердечно-сосудистых осложнений у больных гипотиреозом 62

*Булдигіна Ю.В., Терехова Г.М.,
Болгов М.Ю., Фед'ко Т.В.*

- Патологія щитоподібної залози у пацієнтів з акромегалією: огляд літератури та власні дослідження 67

Муратова Ш.Т.

- Психоэндокринные нарушения у подростков с болезнью Грейвса 71

Експериментальна ендокринологія

Семенко В.В., Сердюк В.М., Савицький І.В.

- Розробка експериментальної аллоксанової моделі цукрового діабету 76

To Practicing Endocrinologists

- What's new in the detection and treatment of dyslipidemia and the prevention of progression of atherosclerosis? 35

Literature Review

*T.V. Sorokman, P.M. Moldovan,
N.O. Popeliuk, O.V. Makarova*

- Genetic basis of endocrine pathology 47

*O.A. Olenovych, N.V. Pashkovska,
V.M. Pashkovskyy, I.V. Boyko*

- Peripheral nervous system damage in hypothyroidism: current view on the problem (literature review) 57

Clinical Thyroidology

*V.S. Vernygorodskyi, N.M. Fetisova,
M.V. Vernygorodskaya*

- Pathogenetic significance of proinflammatory cytokines and dyslipidemia in the development of cardiovascular complications in patients with hypothyroidism 62

*Yu. V. Buldygina, H.M. Terekhova,
M. Yu. Bolgov, T.V. Fedko*

- Thyroid pathology in patients with acromegaly: review of literature and own researches 67

Sh. T. Muratova

- Psycho-endocrine disorders in adolescents with Graves' disease 71

Experimental Endocrinology

V.V. Semenko, V.M. Serdyuk, I.V. Savytskyi

- Development of experimental alloxan model of diabetes mellitus 76

UDC 616.833-036.1-02: 616.44-008.64

DOI: 10.22141/2224-0721.13.4.2017.106652

O.A. Olenovych¹, N.V. Pashkovska¹, V.M. Pashkovskyy¹, I.V. Boyko²¹Higher State Education Institution of Ukraine "Bukovinian State Medical University", Chernivtsi, Ukraine²RME «Chernivtsi Regional Endocrinology Center», Chernivtsi, Ukraine

Peripheral nervous system damage in hypothyroidism: current view on the problem (literature review)

Abstract. The article presents the pathogenesis of polyneuropathy symptoms in patients with hypothyroidism, whereby the approaches to the pathogenetic therapy are substantiated. The article represents the most important aspects of hypothyroid polyneuropathy treatment according to the latest international guidelines and perspective scientific approaches to optimize the medical care for patients with this pathology.

Keywords: hypothyroidism; polyneuropathy; hypothyroid polyneuropathy

Nowadays the thyroid pathology is known to be one of the most widely spread ones in the structure of endocrine diseases [2, 15]. Recently the increase of occurrence of autoimmune thyropathies, mostly followed by the development of hypothyroidism has been observed in Ukraine and other countries. At the same time, the number of surgeries for nodular forms of goiter, tumors, etc., frequently resulting in hypothyroidism, growths as well. The results of epidemiological studies show, that the overall prevalence of manifested hypothyroidism in the population is 0.2–2 %, subclinical one — 7–10 % among women and 2.3 % among men, and morbidity rates are continuously rising every year [15]. There are more than 98 hundred people with this disease, officially registered in Ukraine at the beginning of 2016 [16].

At its early stages the disease is accompanied by a wide range of neurological syndromes, which often dominate in the clinical manifestation of the disease and involve practically all levels of the nervous system [7, 11]. However, despite a long history of detection of interconnections between the thyroid and neurological pathologies, the study of damaging mechanisms for the nervous system in hypothyroidism still remains a topical issue of contemporary neuroendocrinology.

Deficiency of thyroid hormones in the body leads to disorders of water and electrolyte balance, protein, lipid, carbohydrate metabolisms, causing morphological-functional and biochemical changes in various organs and systems [2, 15]. Hypothyroidism is accompanied by disturbed synthesis of neurotransmitters, increased levels of blood lipids, lowering the energy potential of the

cells, the activation of free radical processes, reduction of synthesis of a nitric oxide and endothelial dysfunction, disturbance of blood microcirculation disorders, imbalance of proinflammatory cytokines and adipocytokines, etc. [18]. A dramatic inhibition of energy and anabolic processes, typical for hypothyroidism, promotes an organic damage of the nervous system [22].

Hypothyroid neurological disorders are various and numerous [14, 15]. Marked changes in the peripheral nervous system, typical for hypothyroidism, are implemented in the development of pseudomyotonic and pseudomyasthenic syndromes, radiculopathies, polyneuritis, tunnel neuropathies, and polyneuropathies as well. The latter are found in 18–72 % of patients with hypothyroidism [7, 11, 22], and symptoms of polyneuropathy can develop not only in manifested hypothyroidism, but also in sub-clinical one [30]. However, there is no consensus concerning the pathogenesis of hypothyroid polyneuropathy (PNP), correlation between the degree of its manifestation and hormonal status, state of the neuromuscular system during the compensation of the underlying disease. Thus, indicating a direct connection between the level of thyroid hormones, the degree of hypothyroidism compensation and polyneuropathy symptoms, some researchers believe that all clinical, electroneuromyographic and histopathological changes in patients with hypothyroid polyneuropathy are reversible in case an adequate replacement therapy is initiated [25, 28]. However, according to the other studies, clinical and pathomorphological signs of neuromuscular system disorders remain after the compensation of hypothyroidism [24, 27, 29].

Degenerative, toxic, metabolic, ischemic and mechanical factors, leading to the changes of the connective tissue interstitium, myelin sheath and axial cylinder of the nerve are known to be underlying factors promoting the formation of PNP, particularly hypothyroid one. Hence, the division of polyneuropathy into *axonopathy*, related to the underlying primary damage of the axial cylinders of nerves, and *myelinopathy*, characterized by the disturbance of nerve conduction due to the myelin sheaths loss, is generally accepted. However, during the progression of the disease their combinations usually occur [17].

The development of the PNP in case of hypothyroidism is considered to be related to the mucinous infiltration of the perineurium resulted in nerves compression, as well as to disorders of oxidative processes due to the thyroid hormones insufficiency. Schwann's cells are primarily influenced by the metabolic disorders, and that leads to segmental demyelination. Thus, as morphological studies have demonstrated, glycogen and mucin deposits in Schwann's cells, bulbous thickening of the myelin sheath with mucinous inclusions, segmental demyelination, increased number of demyelinated fibers of a small diameter, reduced number of myelinated fibers of a large diameter are observed in the peripheral nerves [18, 22].

Clinically, hypothyroid PNP is manifested by the pain and paresthesias in the distal parts of the extremities, muscular weakness, seizures, polyneuritic impairment of sensitivity, decrease or loss of tendon reflexes [14, 17]. Such symptoms are amplified to complete immobilization while staying in a cold room or during winter. Severity of PNP clinical manifestation depends on the degree of involvement of motor, sensory and autonomic fibers into the pathological process.

Movement disorders are manifested as muscular weakness, located mainly in the distal areas, mostly in the extensor muscles, accompanied by hypo- or areflexia. In severe cases, patients are unable to stand or walk, hold objects in their hands.

Sensory changes are associated with positive (paresthesia, hyperpathies) and negative symptoms (loss of joint, muscle and tendon proprioception, leading to an imbalance when standing and walking, reduction of skin tactile and pain sensitivity).

Autonomic symptoms appear as sympathalgias, vasoconstrictor, trophic and secretory disorders (burning pain, sweating changes, swelling of the distal areas of the limbs, abnormalities of their color and temperature, sores, muscular changes, nails deformation).

However, the diagnostics of motor, sensory and autonomic symptoms of PNP is generally impeded by multifactorial clinical manifestations of hypothyroidism and involvement of numerous tissues. Thus, the development of hypothyroid myopathy, characterized by permanent muscular weakness (more substantial in muscles of proximal areas of the limbs), prolonged muscular contraction and their relaxation period, by convulsions, muscular hypertrophy and hardening, myalgia, increased mechanical excitability of muscles during percussion, etc., is associated with the limitation and inhibition of move-

ments, mistakenly considered as paresis or paralysis [1]. Moreover, the boundaries of loss and irritation symptoms, the type of sensory disorders, especially in mixed areas, vary in a wide range due to the variability of overlapping of innervation zones by adjacent nerves and variability of autonomous zones, as well as due to the double, triple innervation of certain muscles and skin areas.

In addition to complete neurological examination and selection of typical neurological symptoms a significant role in the differential diagnosis of the type of hypothyroid peripheral nervous system disorder is played by electromyography (EMG). During electromyographic examination the decrease in amplitude and slowing of the speed of an impulse transmission by sensory and motor nerves is recorded, the results of EMG enable to differentiate of the damage of muscle and nerve fibers, detect the degree of nerve fibers' damage, differentiate between axonal (axonopathies) and demyelinating (myelopathy) PNP. Thereby, a slow speed of the transmission of an impulse along the nerve, increased distal latency period, change of F-response, blockage of the transmission and temporary dispersion are usually indicative of the myelin sheath damage, whereas the decrease of impulse level is a sign of axonal degeneration [19].

In doubtful cases nerve biopsy may be very helpful — histological changes in them are absent in case of the progressive muscular degeneration [14, 18].

Furthermore, neurological diagnosis is primarily polysyndromic: it is made according to the prevalence of clinical signs (sensory, motor, autonomic) and the distribution of lesions (symmetrical/asymmetrical, proximal/distal). These statements are important not only in terms of diagnostics, but also for adequate treatment and prognosis.

A particular algorithm to manage patients with hypothyroid PNP doesn't exist. Treatment and rehabilitation of patients with hypothyroidism should be based on hormone replacement therapy in doses, enabling to achieve and maintain euthyroidism [16]. On the other hand, considering those metabolic disorders developed under thyroid hormones insufficiency and causative of the central and peripheral nervous system damage, administration of additional medications normalizing the mentioned changes is substantiated. Therefore, in addition to the treatment of the underlying disease and achievement of hormonal status compensation, two main approaches may be suggested in the therapy of hypothyroid PNP: *pathogenetic therapy* (influence on the mechanisms of nerve fibers damage, stimulation of the regeneration of damaged nerve fibers) and *symptomatic*, targeted on the correction of the symptoms, first of all — on the pain management and patients' quality of life improvement.

Pathogenetic therapy is aimed at slowing of the neuropathy progression and correction of neuropathic deficiency. To achieve this, α -lipoic acid and group B vitamins are used [12, 26].

α -lipoic (thioctic) acid medications create the basis of pathogenetic treatment of PNP [4, 9, 12]. Accumulated in nerve fibers, they inactivate free radicals, block their generation, restore endogenous systems of antiradical protection, thereby providing a powerful antioxidant

effect, and restore a disturbed endoneurial blood flow, normalizes the content of NO (a regulator of the vascular wall relaxation), improves endothelial function, reduces total cholesterol blood level, increases antiatherogenic lipoprotein fraction content [4]. Thioctic acid is a coenzyme to key enzymes of the Krebs cycle that explains its efficacy to optimize the energy metabolism of neurons. This action results in the improvement of nerve conduction by motor and sensory nerve fibers. Moreover, a positive effect of the medication on the liver cells is noticed — reduction of the severity of morphological manifestations of fatty liver and normalization of biochemical parameters [4, 9, 12].

A significant role in pharmacotherapy of PNP belongs to vitamin therapy. Group B vitamins improve the metabolism in the nervous tissue, metabolism of mediators, the transfer of excitation and, as a result, increase the rate of impulse transmission by nerve fibers, as well as implement a moderate analgesic effect, promote the processes of regeneration and remyelination of nerve fibers [5, 6, 13].

As a result of phosphorylation processes thiamin (vitamin B₁) is converted in the body to cocarboxylase, which is a coenzyme of numerous enzymatic reactions and plays an important role in carbohydrate, protein and fat metabolism. Vitamin B₁ is involved in the synthesis of neurotransmitters that modulate the transmission of nerve impulses in the synapses, possesses anticholinesterase activity, hence stipulating neuromuscular conduction; it is one of the components of nucleic acid synthesis and stimulates the plastic and reparative processes in the nervous tissue [10].

Vitamin B₂ (riboflavin) is a catalyst for cell respiration, known to play an important role in redox processes of the nervous system, regulates the metabolism of carbohydrates, proteins, fats, potentiates the effect of pyridoxine and tryptophan, stimulates the regeneration of tissues.

Pyridoxine (vitamin B₆) reduces blood level of cholesterol and lipids, promotes the conversion of folic acid to its active form; it is a coenzyme in the metabolism of amino acids and proteins in the central nervous system cells, in the synthesis of biogenic amines, components of myelin sheath of neurons, neurotransmitters of the central and peripheral nervous system, thus providing a synaptic transmission.

Cyanocobalamin (vitamin B₁₂) provides hematopoietic, erythropoietic, anti-anemic, metabolic action, normalizes blood clotting processes, diversely influences liver function, including hematopoietic one, as well as digestive system, activates the metabolism of carbohydrates and fats, affects the synthesis of RNA, DNA. In addition, vitamin B₁₂ suppresses the abnormal changes in case of degenerative atrophy of the nerve cells, causes resynthesis of myelin, creates the myelin sheath, and thereby provides the restoration of normal nerve fiber structure and functions [5, 10].

Hence, the use of neurotropic group B vitamins can be considered as an important element of pathogenetic therapy of the PNP, promoting the regression of sensitivity disturbances, vegetative symptoms and pain syn-

drome. Since their simultaneous use is essential for the treatment efficacy, the use of combined B vitamins medications is practically expedient and simplifies patients' treatment significantly [5, 6, 13].

In addition to the mentioned drugs pathogenetic therapy of hypothyroid PNP is reasonably contributed by reparants (Actovegin, Solcoseryl) that demonstrate antioxidant, antyhypoxaemic, neurotrophic, neuroprotective action and are widely used in the rehabilitation of patients with various nervous system diseases of the vascular, atrophic, infectious, traumatic and other genesis [8, 20, 23].

In case of motor disorders anticholinesterase agents are used — neostigmine methylsulfate (proserine), ipidacrine (neiromidin), etc. These medications can restore and stimulate the neuromuscular transfer, restore impulse transmission by the peripheral nerves, enhance contractility of the smooth muscles, improve memory and learning ability through stimulation of nervous impulse transmission in the central nervous system as well, specifically moderately stimulating the CNS, provide analgesic effect due to the ability to block sodium permeability of the membranes.

The basis of symptomatic therapy of PNP is preferably related to the correction of the pain syndrome, programmed by the first-line drugs as anticonvulsants of a new generation — gabapentinoids (pregabalin, gabapentin) — soluble amino acids, chemically similar to the endogenous inhibitory neurotransmitter g-aminobutyric acid (GABA), involved in the transmission and modulation of the pain [3, 12].

Gabapentin has a number of biochemical properties that enable its influence on the pathogenesis of chronic neuropathic pain syndrome [3]:

- interaction with $\alpha 2-\delta 2$ -subunits potential-dependent calcium channels, suppression of entry of Ca^{2+} ions into neurons inhibits excessive excitability of cell membranes, reduces sensitization of nociceptors;

- an increase of GABA synthesis stimulates the activity of glutamate decarboxylase, resulting in the enhancement of antinociceptive system activity;

- inhibition of the synthesis of glutamate (stimulating neurotransmitter with exitotoxicity) leads to a decrease in excitability structures nociceptive system and prevents neuronal death;

- modulation of the activity of NMDA (N-methyl-D-aspartate)-receptors affects the processes of «pain memory» formation.

Effective and safe for all types of spontaneous and stimulus-dependent neuropathic pain, gabapentin has flexible circuit of dose titration that provides highly individualized selection of therapy based on clinical characteristics of the patient and his pain syndrome [3].

Pregabalin is a modern anticonvulsant that has shown its efficacy regarding any type of neuropathic pain, fibromyalgia, seizures with a high analgesic activity and a positive impact on concomitant emotional-depression manifestations [21].

Treatment algorithm for neuropathic pain in PNP also includes tricyclic antidepressants (eg, amitriptyline), selective serotonin reuptake inhibitors (duloxetine,

venlafaxine), etc. [12], but their psychotropic side effects and cholinolytic action significantly limit their administration in patients with hypothyroidism.

Non-pharmacologic strategies for management of painful neuropathy include physiotherapeutic techniques (acupuncture, transcutaneous electrical stimulation, high-wave external muscle stimulation, etc.).

Thus, neuromuscular disorders in case of thyropathies, including hypothyroidism, are known as polymorphic and still create significant diagnostic and therapeutic challenges. The processes of demyelination with secondary axonal damage, resulted from metabolic and bioenergetic disorders, initiated by thyroid hormones insufficiency, underlie the development of dysmetabolic polyneuropathy in hypothyroidism. Concerning this, early diagnosis, adequate replacement therapy and achievement of hormonal status compensation remain the major concerns to prevent and postpone the progression of hypothyroid polyneuropathy. However, pathophysiological targeted therapy is a meaningful approach in management strategies for neuropathic changes, that emphasizes the importance of further study of pathogenesis, clinical and neurophysiological manifestations of hypothyroid polyneuropathy, particularly during the compensation period of the underlying endocrine disease, and develops algorithms for differential treatment.

Conflicts of interests. Authors declare the absence of any conflicts of interests that might be construed to influence the results or interpretation of their manuscript.

References

1. Антелава О.А. Дифференциальная диагностика мышечной слабости / О.А. Антелава, Л.Ф. Касаткина., Г.Т. Гуркина и др. // Рос. мед. журнал. — 2004. — № 14. — С. 15-18.
2. Балаболкин М.И. Дифференциальная диагностика и лечение эндокринных заболеваний: Руководство / М.И. Балаболкин, Е.М. Клебанова, В.М. Креминская. — М.: Медицинское информационное агентство, 2008. — 752 с.
3. Баринов А.Н. Алгоритмы диагностики и лечения невропатической боли при поражении периферической нервной системы / А.Н. Баринов, Л.Т. Ахмеджанова, К.А. Махинов // Рос. мед. журнал. — 2016. — № 3. — С. 154-162.
4. Больщова О.В. Застосування альфа-ліпоєвої кислоти в комплексному лікуванні порушень ліпідного обміну у дітей та підлітків, хворих на вроджений гіпотиреоз / О.В. Больщова, В.А. Музь, Т.М. Малиновська // Ліки України. — 2011. — № 10(156). — С. 84-88.
5. Вернигородский В.С. Нейровитан в комплексном лечении и реабилитации больных гипотиреозом / В.С. Вернигородский, М.В. Власенко, Н.М. Фетисова // Проблемы эндокринной патологии. — 2011. — № 2. — С. 36-40.
6. Власенко М.В. Якість життя хворих на цукровий діабет 2 типу з метаболічною нейропатією на тлі патогенетичної терапії / М.В. Власенко // Міжнародний ендокринологічний журнал. — 2013. — № 1(49). — С. 1-4.
7. Григорова И.А. Особенности формирования гипотиреоидных полиневропатий / И.А. Григорова, Е.Л. Товажнянская // Укр. неврол. журнал. — 2008. — № 2. — С. 67-72.
8. Кадыков А.С. Нейрометаболическая терапия больных с заболеваниями нервной системы. Возможности применения Актовегина / А.С. Кадыков, Н.В. Шахпаронова // Рос. мед. журнал. — 2008. — № 26. — С. 1722.
9. Корпачев В.В. Лекарственные формы тиоктовой кислоты / В.В. Корпачев, М.И. Борщевская // Проблемы эндокринной патологии. — 2006. — № 1. — С. 1-11.
10. Луцкий И.С. Сравнение эффективности инъекционных и таблетированных форм витамина группы В при лечении полинейропатий / И.С. Луцкий, Я.А. Гончарова, С.К. Евтушенко и др. // Міжнародний неврологічний журнал. — 2009. — № 1 (23). — С. 90-95.
11. Макар Р.Д. Гіпотиреоз: концептуальні аспекти крізь призму часу / Р.Д. Макар, М.В. Сандрурська // Міжнародний ендокринологічний журнал. — 2009. — № 1(19). — С. 124-131.
12. Маньковский Н.Б. Оптимизация медикаментозного лечения диабетической нейропатии / Н.Б. Маньковский // Здоров'я України. — 2007. — № 8. — С. 58.
13. Матвієнко Ю.О. Запальні деміелінізуючі полінейропатії у загальномедичній практиці / Ю.О. Матвієнко, Н.Л. Боженко. — К.: ТОВ «Люди в білому», 2014. — 36 с.
14. Мументалер М. Дифференциальный диагноз в неврологии: руководство по оценке, классификации и дифференциальной диагностике неврологических симптомов / М. Мументалер, К. Бассетти, К. Дэтвайлер. — М.: МЕДпресс-информ, 2010. — 359 с.
15. Паньків В.І. Синдром гіпотиреозу // Міжнародний ендокринологічний журнал. — 2012. — № 5 (45). — С. 174-191.
16. Пашковська Н.В. Лікування гіпотиреозу згідно із сучасними клінічними настановами / Н.В. Пашковська // Міжнародний ендокринологічний журнал. — 2016. — № 6(78). — С. 48-58.
17. Силантьев К. Классическая неврология: руководство по периферической нервной системе и хроническим болевым синдромам / К. Силантьев. — Волгоград: Панорама, 2006. — 400 с.
18. Скрипник Н.В. Патогенетичні основи неврологічних порушень у хворих на гіпотиреоз та їх корекція / Н.В. Скрипник, В.А. Гриб, Н.І. Кубарич, Т.С. Вацеба // Практикуючий лікар. — 2012. — № 4. — С. 55-60.
19. Соловьев А.А. Электронейромиографические параметры нервов верхних конечностей у больных хроническим гипотиреозом / А.А. Соловьев, С.А. Маризин // Сб. статей: Эндокринологической службе Удмуртии 50 лет. — Ижевск, 2005. — С. 182-188.
20. Строков И.А. Актовегин в лечении заболеваний нервной системы / И.А. Строков, К.И. Строков, Е.Е. Даева, Ж.А. Афонина // Рос. мед. журнал. — 2008. — № 12. — С. 1665.
21. Строков И.А. Прегабалин в терапии болевой диабетической полиневропатии / И.А. Строков, А.С. Фокина, В.А. Головачева // Рос. мед. журнал. — 2013. — № 9. — С. 422.
22. Товажнянская Е.Л. Неврологические осложнения первичного гипотиреоза различного генеза / Е.Л. Товажнянская // Международный медицинский журнал. — 2013. — № 1. — С. 15-19.
23. Шишкова В.Н. Перспективы применения препарата Актовегин у пациентов с метаболическим синдромом и предиабетом. Современные представления о нарушениях углеводного обмена / В.Н. Шишкова // Рос. мед. журнал. — 2007. — № 27. — С. 2066.

24. Beghi E. Hypothyroidism and polyneuropathy / E. Beghi, M.L. Delodovici, G. Bogliun et al. // J. Neurol. Neurosurg. Psychiatry. — 1989. — Vol. 52, № 12. — P. 1420-1423.
25. El-Salem K. Neurophysiological changes in neurologically asymptomatic hypothyroid patients: a prospective cohort study / K. El-Salem, F. Ammani // J. Clin. Neurophysiol. — 2006. — № 23(6). — P. 568-572.
26. James C. Watson. Peripheral neuropathy: a practical approach to diagnosis and symptom management / James C. Watson, P. James, B. Dyck // Mayo Clin. Proc. — 2015. — Vol. 90, № 7. — P. 940-951.
27. Kececi H. Hormone replacement therapy in hypothyroidism and nerve conduction study / H. Kececi, Y. Degirmenci // Neurophysiol. Clin. — 2006. — Vol. 36(2). — P. 79-83.
28. Machicao F. Плейотропные нейропротекторные и метаболические эффекты Актовегина / F. Machicao, D.F. Muresanu, H. Hundsberger et al. // Therapia. — 2013. — № 2(77). — С. 2-7.
29. Mahadule A.A. Motor conduction parameters in recently diagnosed and untreated hypothyroidism / A.A. Mahadule, P.S. Jadhao, M.S. Phatak // Ann. Neurosci. — 2015. — Vol. 22(1). — P. 6-10.
30. Penza P. Painful neuropathy in subclinical hypothyroidism: clinical and neuropathological recovery after hormone replacement therapy / P. Penza, R. Lombardi, F. Camozzi et al. // Neurol. Sci. — 2009. — Vol. 30(2). — P. 149-151.

Received 12.06.2017

Оленович О.А.¹, Пашковська Н.В.¹, Пашковський В.М.¹, Бойко І.В.²¹Вищий державний навчальний заклад України «Буковинський державний медичний університет», м. Чернівці, Україна²ОКУ «Чернівецький обласний ендокринологічний центр», м. Чернівці, Україна

Ураження периферичної нервової системи при гіпотиреозі: сучасний погляд на проблему (огляд літератури)

Резюме. У статті наведені дані про патогенез симптомів полінейропатії у хворих на гіпотиреоз, на підставі чого обґрунтовані шляхи патогенетичної терапії. Наведені найбільш важливі аспекти лікування полінейропатії на тлі гіпотиреозу згідно з новітніми міжнародними реко-

мендаціями, зазначені перспективні наукові напрямки для оптимізації надання допомоги пацієнтам із цією патологією.

Ключові слова: гіпотиреоз; полінейропатія; гіпотиреоїдність; полінейропатія

Оленович О.А.¹, Пашковская Н.В.¹, Пашковский В.М.¹, Бойко И.В.²¹Высшее государственное учебное заведение Украины «Буковинский государственный медицинский университет», г. Черновцы, Украина²ОКУ «Черновицкий областной эндокринологический центр», г. Черновцы, Украина

Поражение периферической нервной системы при гипотиреозе: современный взгляд на проблему (обзор литературы)

Резюме. В статье представлены данные относительно патогенеза симптомов полинейропатии у больных гипотиреозом, на основании чего обоснованы пути патогенетической терапии. Приведены наиболее важные аспекты лечения полинейропатии на фоне гипотиреоза согласно

современным международным рекомендациям, указаны перспективные научные направления для оптимизации оказания помощи пациентам с этой патологией.

Ключевые слова: гипотиреоз; полинейропатия; гипотиреоидная полинейропатия