

**МІНІСТЕРСТВО ОХОРОНИ ЗДОРОВ'Я УКРАЇНИ
ВИЩИЙ ДЕРЖАВНИЙ НАВЧАЛЬНИЙ ЗАКЛАД УКРАЇНИ
«БУКОВИНСЬКИЙ ДЕРЖАВНИЙ МЕДИЧНИЙ УНІВЕРСИТЕТ»**



МАТЕРІАЛИ

101 – ї

підсумкової наукової конференції

професорсько-викладацького персоналу

Вищого державного навчального закладу України

«БУКОВИНСЬКИЙ ДЕРЖАВНИЙ МЕДИЧНИЙ УНІВЕРСИТЕТ»

10, 12, 17 лютого 2020 року

Чернівці – 2020

УДК 001:378.12(477.85)
ББК 72:74.58
М 34

Матеріали 101 – ї підсумкової наукової конференції професорсько-викладацького персоналу вищого державного навчального закладу України «Буковинський державний медичний університет» (м. Чернівці, 10, 12, 17 лютого 2020 р.) – Чернівці: Медуніверситет, 2020. – 488 с. іл.

ББК 72:74.58

У збірнику представлені матеріали 101 – ї підсумкової наукової конференції професорсько-викладацького персоналу вищого державного навчального закладу України «Буковинський державний медичний університет» (м.Чернівці, 10, 12, 17 лютого 2020 р.) із стилістикою та орфографією у авторській редакції. Публікації присвячені актуальним проблемам фундаментальної, теоретичної та клінічної медицини.

Загальна редакція: професор Бойчук Т.М., професор Іващук О.І.,
доцент Безрук В.В.

Наукові рецензенти:

професор Братенко М.К.
професор Булик Р.Є.
професор Гринчук Ф.В.
професор Давиденко І.С.
професор Дейнека С.Є.
професор Денисенко О.І.
професор Заморський І.І.
професор Колоскова О.К.
професор Коновчук В.М.
професор Пенішкевич Я.І.
професор Сидорчук Л.П.
професор Слободян О.М.
професор Ткачук С.С.
професор Тодоріко Л.Д.
професор Юзько О.М.
професор Годованець О.І.

ISBN 978-966-697-843-4

© Буковинський державний медичний
університет, 2020



Kryvetskyi I.V.

**PECULIARITIES OF MORPHOGENESIS AND TOPOGRAPHY OF THE THORACIC
SPINE OF THE SPINAL COLUMN IN THE PRENATAL PERIOD OF HUMAN
ONTOGENESIS**

*Department of Anatomy, Topographic Anatomy and Operative Surgery
Higher State Educational Establishment of Ukraine
«Bukovinian State Medical University»*

The urgency of the work is explained by the necessity of a complex study of the development peculiarities, topography formation of structures of the thoracic spine of the spinal column and dynamics of their syntopic correlation in the prenatal period of ontogenesis and in the newborns, that is of great significance for elucidation of the morphological preconditions and time of the possible origin of the congenital spinal defects with the object of the development of new, more rational methods of surgical interventions in this area, elaboration of new stabilization technologies and spinal column correction at disabling deformities of the spine in children and adolescents.

The aim is at the ascertainment of chronological sequence of the development and formation of the topography structures of the section of the thoracic part of the spinal column in the early period of human ontogenesis. The topographic and anatomical features of the relationships between the structures of the thoracic part of the spinal column from the moment of their laying to birth, dynamics of their formation and growth taking into account morphogenesis of the adjacent structures are established. With the help of the adequate morphological methods, investigation of morphogenesis and dynamics of spatial-time relationships of the thoracic spine of the spinal column of a person, their connections during the fetal period of the development and in the newborns from the point of view of the topographic-anatomical approach to embryogenesis problems was carried out. The features of the blood supply and venous outflow of the spine are ascertained. Critical periods, morphological preconditions and time of the possible origin of some innate defects of the spinal column were established. On the basis of the obtained results, the problem of prenatal diagnostics of the innate malformations of the thoracic part of the spinal column was solved.

The thoracic vertebrae laying occurs in the germs of 7.0-9.0 mm CRL by forming the condensation of sclerotome cells round the chord and the nervous tube, from which mesenchymal thoracic vertebrae are formed. The vertebral bodies are formed from the cranial and caudal parts of two adjacent sclerotome masses. Intersegmental arteries remain on the level of the vertebral bodies, and the spinal nerves lie between thoracic vertebrae. In the germs of 10.0-12.5 mm CRL the arches of the vertebrae move away from the bodies perpendicularly in the dorsal direction.

The formation of articular and transverse processes begin. At this early stage of the development there are no joints in the spinal column of the germs, the spinal canal forming begins. Bodies are clearly defined from the thoracic vertebrae, and in the lumbar and sacral vertebrae only arches are clearly visible and closely spaced bodies. The vertebral bodies at this stage are well differentiated. All of them have the same, primitive, quadrilateral body shape and are separated from each other by a layer of mesenchyma. The layers correspond to the future intervertebral discs.

Kryvetskyi V.V.

**PECULIARITIES OF MORPHOGENESIS AND TOPOGRAPHY OF THE THORACIC
PART OF THE VERTEBRAL COLUMN IN THE PRENATAL PERIOD OF HUMAN
ONTOGENESIS**

*M.G. Turkevich Department of Human Anatomy
Higher State Educational Establishment of Ukraine
„Bukovinian State Medical University”*

The urgency of the work is explained by the necessity of a complex study of the development peculiarities, topography formation of structures of the thoracic spine of the spinal column and dynamics of their syntopic correlation in the prenatal period of ontogenesis and in the newborns, that is of great significance for elucidation of the morphological preconditions and time