



colloquium-journal

ISSN 2520-6990

*Międzynarodowe czasopismo naukowe*



INDEX COPERNICUS  
INTERNATIONAL

Google  
scholar

issuu™

calaméo



SlideShare




НАУЧНАЯ ЭЛЕКТРОННАЯ  
БИБЛИОТЕКА

**LIBRARY.RU**

Jurisprudence  
Medical sciences  
Chemical sciences  
Economic sciences  
Biological sciences  
Philological sciences  
Pedagogical sciences

№33(156) 2022



ISSN 2520-6990

ISSN 2520-2480

Colloquium-journal №33 (156), 2022

Część 1

(Warszawa, Polska)

Redaktor naczelny - **Paweł Nowak**  
**Ewa Kowalczyk**

Rada naukowa

- **Dorota Dobija** - profesor i rachunkowości i zarządzania na uniwersytecie Koźmińskiego
- **Jemielniak Dariusz** - profesor dyrektor centrum naukowo-badawczego w zakresie organizacji i miejsc pracy, kierownik katedry zarządzania Międzynarodowego w Ku.
- **Mateusz Jabłoński** - politechnika Krakowska im. Tadeusza Kościuszki.
- **Henryka Danuta Stryczewska** – profesor, dziekan wydziału elektrotechniki i informatyki Politechniki Lubelskiej.
- **Bulakh Iryna Valerievna** - profesor nadzwyczajny w katedrze projektowania środowiska architektonicznego, Kijowski narodowy Uniwersytet budownictwa i architektury.
- **Leontiev Rudolf Georgievich** - doktor nauk ekonomicznych, profesor wyższej komisji atestacyjnej, główny naukowiec federalnego centrum badawczego chabarowska, dalekowschodni oddział rosyjskiej akademii nauk
- **Serebrennikova Anna Valerievna** - doktor prawa, profesor wydziału prawa karnego i kryminologii uniwersytetu Moskiewskiego M.V. Lomonosova, Rosja
- **Skopa Vitaliy Aleksandrovich** - doktor nauk historycznych, kierownik katedry filozofii i kulturoznawstwa
- **Pogrebnaya Yana Vsevolodovna** - doktor filologii, profesor nadzwyczajny, stawropolski państwowy Instytut pedagogiczny
- **Fanil Timeryanowicz Kuzbekov** - kandydat nauk historycznych, doktor nauk filologicznych. profesor, wydział Dziennikarstwa, Bashgosuniversitet
- **Aliyev Zakir Hussein oglu** - doctor of agricultural sciences, associate professor, professor of RAE academician RAPVHN and MAEP
- **Kanivets Alexander Vasilievich** - kandydat nauk technicznych, profesor nadzwyczajny Wydział Agrotechnologii i Transportu Drogowego, Państwowy Uniwersytet Rolniczy w Połtawie
- **Yavorska-Vitkovska Monika** - doktor edukacji, szkoła Kuyavsky-Pomorski w Bydgoszczy, dziekan nauk o filozofii i biologii; doktor edukacji, profesor
- **Chernyak Lev Pavlovich** - doktor nauk technicznych, profesor, katedra technologii chemicznej materiałów kompozytowych narodowy uniwersytet techniczny Ukrainy „Politechnika w Kijowie”
- **Vorona-Slivinskaya Lyubov Grigoryevna** - doktor nauk ekonomicznych, profesor, St. Petersburg University of Management Technologia i ekonomia
- **Voskresenskaya Elena Vladimirovna** doktor prawa, kierownik Katedry Prawa Cywilnego i Ochrony Własności Intelektualnej w dziedzinie techniki, Politechnika im. Piotra Wielkiego w Sankt Petersburgu
- **Tengiz Magradze** - doktor filozofii w dziedzinie energetyki i elektrotechniki, Georgian Technical University, Tbilisi, Gruzja
- **Usta-Azizova Dilnoza Ahrarova** - kandydat nauk pedagogicznych, profesor nadzwyczajny, Tashkent Pediatric Medical Institute, Uzbekistan
- **Oktay Salamov** - doktor filozofii w dziedzinie fizyki, honorowy doktor-profesor Międzynarodowej Akademii Ekoenergii, docent Wydziału Ekologii Azerbejdżańskiego Uniwersytetu Architektury i Budownictwa
- **Karakulov Fedor Andreevich** – researcher of the Department of Hydraulic Engineering and Hydraulics, federal state budgetary scientific institution "all-Russian research Institute of hydraulic Engineering and Melioration named after A. N. Kostyakov", Russia.

Google Scholar SlideShare



INDEX  
INTERNATIONAL



COPERNICUS



НАУЧНАЯ ЭЛЕКТРОННАЯ  
БИБЛИОТЕКА  
LIBRARY.RU

«Colloquium-journal»

Wydawca «Interdruk» Poland, Warszawa

Annopol 4, 03-236

E-mail: [info@colloquium-journal.org](mailto:info@colloquium-journal.org)

<http://www.colloquium-journal.org/>

# CONTENTS

## MEDICAL SCIENCES

<b>Antoniv A.A., Menzak Yu.T.</b> THE MECHANISMS OF INTERACTION OF NON-ALCOHOLIC FATTY LIVER DISEASE AND CHRONIC KIDNEY DISEASE .....	4
<b>Антонів А.А., Мензак Ю.Т.</b> МЕХАНІЗМИ ВЗАЄМОБ'ЯЖЕННЯ НЕАЛКОГОЛЬНОЇ ЖИРОВОЇ ХВОРОБИ ПЕЧІНКИ ТА ХРОНІЧНОЇ ХВОРОБИ НИРОК .....	4
<b>Antoniv A.A., Shakula I.V.</b> INTENSITY OF LIVER FIBROSIS IN PATIENTS WITH NON-ALCOHOLIC STEATOHEPATITIS ON THE BACKGROUND OF OBESITY WITH COMORBIDITY WITH CHRONIC KIDNEY DISEASE I-III ST. ....	7
<b>Антонів А.А., Шакула І.В.</b> ІНТЕНСИВНІСТЬ ФІБРОЗУ ПЕЧІНКИ У ХВОРИХ НА НЕАЛКОГОЛЬНИЙ СТЕАТОГЕПАТИТ НА ТЛІ ОЖИРІННЯ ЗА КОМОРБІДНОСТІ З ХРОНІЧНОЮ ХВОРОБОЮ НИРОК I-III СТАДІЇ .....	7
<b>Antoniv A.A., Verbych V.Yu.</b> ENDOTHELIAL DYSFUNCTION IN THE PATHOGENESIS OF NONALCOHOLIC STEATOHEPATITIS IN PATIENTS WITH CHRONIC KIDNEY DISEASE .....	10
<b>Антонів А.А., Вербич В.Ю.</b> ЕНДОТЕЛІАЛЬНА ДИСФУНКЦІЯ В ПАТОГЕНЕЗІ НЕАЛКОГОЛЬНОГО СТЕАТОГЕПАТИТУ У ХВОРИХ НА ХРОНІЧНУ ХВОРОБУ НИРОК.....	10
<b>Antoniv A.A., Komendant V.A.</b> PATHOGENETIC FEATURES OF NON-ALCOHOLIC FATTY LIVER DISEASE AND CHRONIC KIDNEY DISEASE .....	13
<b>Антонів А.А., Комендант В.А.</b> ПАТОГЕНЕТИЧНІ ОСОБЛИВОСТІ ПЕРЕБІГУ НЕАЛКОГОЛЬНОЇ ЖИРОВОЇ ХВОРОБИ ПЕЧІНКИ ТА ХРОНІЧНОЇ ХВОРОБИ НИРОК.....	13
<b>Antoniv A.A., Tseberskyi K.A.</b> THE TREATMENT OF NON-ALCOHOLIC STEATOHEPATITIS IN COMORBID COURSE WITH CHRONIC KIDNEY DISEASE (CHRONIC PYELONEPHRITIS) .....	17
<b>Антонів А.А., Цеберський К.А.</b> ЛІКУВАННЯ НЕАЛКОГОЛЬНОГО СТЕАТОГЕПАТИТУ ЗА КОМОРБІДНОГО ПЕРЕБІГУ З ХРОНІЧНОЮ ХВОРОБОЮ НИРОК (ХРОНІЧНИЙ ПІЕЛОНЕФРИТ) .....	17
<b>Ivanova L.A., Drutsul-Melnyk N.V., Komar O., Kropatnytska Ya., Horodynska Yu., Honchar A.</b> FEATURES OF THE COURSE AND COMPLICATIONS OF COVID-19 IN CHILDREN (LITERATURE REVIEW).....	22
<b>Sokolenko M.O., Anohina D.D.</b> RISK OF CHOLERA EPIDEMIC DEVELOPMENT IN UKRAINE: CURRENT STATE OF THE PROBLEM.....	24
<b>Соколенко М.О., Анохіна Д.Д.</b> РИЗИК РОЗВИТКУ ЕПІДЕМІЇ ХОЛЕРИ В УКРАЇНІ: СУЧАСНИЙ СТАН ПРОБЛЕМИ .....	24
<b>Sokolenko M.O., Khovanets K.R.</b> BRIEF INFORMATION ON THE DISTRIBUTION AND CLINICAL MANIFESTATIONS OF MENINGITIS AMONG UKRAINIAN CHILDREN .....	26
<b>Соколенко М.О., Хованець К.Р.</b> СТИСЛІ ВІДОМОСТІ ПРО ПОШИРЕННЯ ТА КЛІНІЧНІ ПРОЯВИ МЕНІНГІТУ СЕРЕД ДІТЕЙ УКРАЇНИ .....	26
<b>PEDAGOGICAL SCIENCES</b>	
<b>Малишевська О.С., Гречух Л.С., Пасько В.А., Тимків А. І.</b> ВПРОВАДЖЕННЯ КОНЦЕПЦІЇ СТАЛОГО РОЗВИТКУ У ВИЩІЙ МЕДИЧНІЙ ШКОЛІ.....	29
<b>Malyshevska O.S., Hrechuch L.S., Pasko V.A., Tymkiv A.I.</b> IMPLEMENTATION OF THE CONCEPT OF SUSTAINABLE DEVELOPMENT IN THE HIGHER MEDICAL SCHOOL .....	29
<b>Antoniv A.A., Kushnir A.R.</b> THE INFLUENCE OF MODERN TECHNOLOGIES ON THE QUALITY OF PROFESSIONAL TRAINING OF STUDENTS OF MEDICAL EDUCATIONAL INSTITUTIONS.....	33
<b>Антонів А.А., Кушнір А.Р.</b> ВПЛИВ СУЧАСНИХ ТЕХНОЛОГІЙ НА ЯКІСТЬ ПРОФЕСІЙНОГО ПІДГОТОВКИ СТУДЕНТІВ МЕДИЧНИХ ОСВІТНІХ ЗАКЛАДІВ .....	33
<b>Білявська Т. М.</b> ГРУПОВА ФОРМА ОРГАНІЗАЦІЇ НАВЧАЛЬНОЇ ДІЯЛЬНОСТІ В ПРОЦЕСІ МОВНОЇ ПІДГОТОВКИ МАЙБУТНІХ УЧИТЕЛІВ ПОЧАТКОВОЇ ШКОЛИ .....	35
<b>Biliavska T. M.</b> GROUP FORM OF ORGANIZING EDUCATIONAL ACTIVITIES IN THE PROCESS OF LANGUAGE TRAINING OF FUTURE PRIMARY SCHOOL TEACHERS.....	35

Antoniv A.A.,

Verbych V.Yu.

Bukovynian State Medical University

**ENDOTHELIAL DYSFUNCTION IN THE PATHOGENESIS OF NONALCOHOLIC  
STEATOHEPATITIS IN PATIENTS WITH CHRONIC KIDNEY DISEASE**

Антонів А.А.,

Вербич В.Ю.

ВДНЗ України «Буковинський державний медичний університет»

**ЕНДОТЕЛІАЛЬНА ДИСФУНКЦІЯ В ПАТОГЕНЕЗІ НЕАЛКОГОЛЬНОГО СТЕАТОГЕПАТИТУ  
У ХВОРИХ НА ХРОНІЧНУ ХВОРОБУ НИРОК****Анотація.**

У статті наведено теоретичне узагальнення результатів дослідження ендотеліальної дисфункції за коморбідного перебігу неалкогольного стеатогепатиту з ожирінням та хронічною хворобою нирок I-II стадії, що супроводжується глибшим ліпідний дисбаланс, гіперхолестеролемія, у т.ч. у складі ліпопротеїдів низької густини, зниження вмісту ліпопротеїдів високої густини, зростання індексу атерогенності, а також гіперпродукція NO ендотелієм та лімфоцитами із прогресуючим ушкодженням ендотелію.

**Abstract.**

The article presents a theoretical generalization of the results of the study of endothelial dysfunction in the comorbid course of non-alcoholic steatohepatitis with obesity and chronic kidney disease stage I-II, accompanied by a deeper lipid imbalance, hypercholesterolemia, including in the composition of low-density lipoproteins, decrease in the content of high-density lipoproteins, increase in the atherogenicity index, as well as hyperproduction of NO by endothelium and lymphocytes with progressive endothelial damage.

**Keywords:** non-alcoholic steatohepatitis, chronic kidney disease, endothelial dysfunction.

**Ключові слова:** неалкогольний стеатогепатит, хронічна хвороба нирок, ендотеліальна дисфункція.

**Introduction.** The comorbidity of non-alcoholic steatohepatitis (NASH) and chronic kidney disease (CKD) on the background of obesity is often recently drawn to the attention of both practitioners and researchers [1, 2]. Schematically, the development of NASH can be presented in several stages: fatty infiltration of the liver, oxidative stress, mitochondrial dysfunction, TNF / endotoxin-mediated injury, aseptic inflammation, diffused liver fibrosis, development of liver-cellular insufficiency (LCI) [1,2,3]. The first place among the causes of the development of NASH is insulin resistance syndrome. NASH most often occurs in obesity (20-81%). The prevalence of NASH in the world is 10% (600 million people) [2,4,5]. In the last 5 years in Ukraine, the incidence of steatohepatitis has increased by 76.6%. In the 12-40% of patients with liver steatosis during 8-13 years, NASH develops with early liver fibrosis (LF).

Chronic kidney disease (CKD) is an important problem in Ukraine and the world today, and the incidence rate has increased by 17% in recent years.

The frequency of occurrence of NASH in patients with CKD is unknown. The mechanisms of their joint development are described in isolated works, which were conducted mainly in the experiments. Despite the fact that among various pathological processes in the internal organs that occur in the background of a metabolic syndrome - NASH is an extremely common disease, and quite often it occurs in patients with CKD, so far, this comorbidity remains a significant problem of the present and needs to be sufficiently studied.

**The aim of the study** – to find out the features of the lipid blood spectrum and the development of endothelial dysfunction in non-alcoholic steatohepatitis in patients with obesity of the 1st degree and chronic kidney disease of the 1st and 2nd stage.

**Material and methods of research.** 135 patients with non-alcoholic steatohepatitis (NASH) with comorbid obesity I degree and chronic kidney disease (CKD) of the 1st and 2nd stage, were examined. Patients were divided into 2 groups: of which 52 patients with non-alcoholic steatohepatitis with obesity I degree (group 1), 53 patients with non-alcoholic steatohepatitis with comorbid obesity of the 1st degree and chronic kidney disease of the I-II stage (chronic uncomplicated pyelonephritis with latent phase in subsiding exacerbation phase) (group 2). The control group consisted of 30 practically healthy persons of the corresponding age and sex. The average age of patients was  $(45.8 \pm 3.81)$  years, men were 48, and women 57 persons.

The functional state of the endothelium was studied by the content of stable metabolites of nitrogen monoxide (NO) (nitrites, nitrates) In the blood by L.C.Green et al. The number of desquamated endothelial cells in the blood was determined by the method of J.Hladovec in the modification of N.N. Petrishev et al. The statistical analysis was performed using parametric and non-parametric criteria (Student, Pearson) on RS AMD Athlon 64 using Statistica 5.1 software (StatSoft, Inc., USA) and SPSS 10.0.5. Standard Version.

The diagnosis of NASH was established in accordance with the unified clinical protocol, approved by the order of the Ministry of Health of Ukraine No. 826 from 06.11.2014, in the presence of criteria for the exclusion

of chronic diffuse liver disease of the viral, hereditary, autoimmune or medicinal genesis as causes of cholestatic or cytolytic syndromes, as well as the results of the USG survey. Diagnosis and treatment of CKD were performed according to the recommendations of the clinical guidelines of the State Institute "Institute of Nephrology, NAMS of Ukraine" (2012). The analysis of clinical manifestations of NASH and CKD of the 1st and 2nd stage, biochemical, laboratory parameters of the functional state of the liver, kidneys, endothelium, ultrasonographic data was studied in dynamics after 30, 90 days of treatment, as well as 3 months after treatment.

The statistical analysis of the results was carried out in accordance with the type of research carried out and the types of numerical data that were obtained. Distribution normality was verified using Liliefors, Shapiro-Wilk tests and the direct visual evaluation of eigenvalues distribution histograms. Quantitative indices having a normal distribution are represented as mean (M)  $\pm$  standard deviation (S). Discrete values are presented in the form of absolute and relative frequencies (percentage of observations to the total number of surveyed). For comparisons of data that had a normal

distribution pattern, parametric tests were used to estimate the Student's t-criterion, Fisher's F-criterion. In the case of abnormal distribution, the median test, Mann-Whitney Rank U-Score, and Wilcoxon's T-criterion (in the case of dependent groups) were used for multiple comparison. Statistica for Windows version 8.0 (Stat Soft inc., USA), Microsoft Excel 2007 (Microsoft, USA) software packages were used for statistical and graphical analysis of the obtained results.

#### Results of the research and their discussion.

Analysis of the lipid profile of the blood in patients with NASH and obesity showed a number of changes that differed depending on the presence of CKD (table 1). Indicators of concentration in blood of total lipids in patients of 1st and 2nd groups exceeded the norm by 26.4% and 34.2%, respectively, with a statistically significant difference between the groups ( $p < 0.05$ ). The content of total cholesterol in blood indicated that it increased by 37.4 and 46.7 ( $p < 0.05$ ) compared with PHPs in patients of 1st and 2nd groups. Changes in the concentration of TG in the form of a significant increase (respectively, 2.2 and 2.0 times ( $p < 0.05$ )) were recorded in the 1st and 2nd groups of patients.

Table 1

**Indicators of lipid spectrum of blood, glycemia and endothelial dysfunction in patients with non-alcoholic stethohepatitis, obesity of the I-II degree and with comorbidity with chronic kidney disease of the I-II stage (M  $\pm$  m)**

Indicators, units measurement	Groups of patients surveyed		
	PHP	Group 1 NASH+Obesity	Group 2 NASH with CKD + Obesity
Total Cholesterol, mmol / l	4,72 $\pm$ 0,11	6,89 $\pm$ 0,38*	6,93 $\pm$ 0,39*/**
LDL, mmol / l	2,54 $\pm$ 0,02	4,05 $\pm$ 0,022 *	4,58 $\pm$ 0,04*/**
HDL, mmol / l	1,28 $\pm$ 0,05	0,72 $\pm$ 0,02 *	0,76 $\pm$ 0,04 */**
TG, mmol / l	1,47 $\pm$ 0,03	2,42 $\pm$ 0,03*	3,19 $\pm$ 0,07 */**
NO IN BLOOD, mmol / l	15,32 $\pm$ 1,225	30,49 $\pm$ 1,318 *	40,51 $\pm$ 1,173 */**
ET-1, pmol / l	6,17 $\pm$ 0,854	11,25 $\pm$ 0,457 *	18,83 $\pm$ 0,559 */**
DEC x104/L	3,03 $\pm$ 0,204	3,87 $\pm$ 0,123 *	5,80 $\pm$ 0,127 */**

Notes: \* - changes are probable in comparison with the index in PHP ( $P < 0,05$ );

\*\* - changes are probable when comparing the indices in patients with NASH ( $P < 0.05$ );

That is, the content in TG in the blood in the comorbid flow of NASH with CKD and obesity were significantly lower than in patients with NASH and obesity.

The study of blood concentrations of proatherogenic lipoprotein fractions indicated a number of changes: the concentration of LDL in the patients of the 1st group was 1.5 times higher than the control group ( $p < 0.05$ ), and in patients of the 2nd group LDL increased in 1.7 times ( $p < 0.05$ ). It is also necessary to point out that with the increase in the activity of cytotoxicity, the content of the cholesterol and LDL in the blood in NASH with comorbidity with CKD and obesity - increased, which may be an important prognostic factor in the progression of atherosclerosis in these patients. Concentration in blood of antiatherogenic lipoproteins - HDL in patients of both groups was significantly lower in comparison with control: in patients of the 1st group - in 1,5 times ( $p < 0,05$ ), in 2nd group - 1,7 times ( $p < 0.05$ ). As can be seen from the results of the study, the maximum suppression of HDL synthesis (Table 1) was observed in patients of the 2nd group, indicating a

minimum level of protection of endothelial vessels from free radical aggression and atherogenic fractions of blood lipids. The result of these changes was a significant increase in the index of atherogenicity in patients of both groups of observation: the 1st group - 2.2 times, the 2nd group - 2.0 times with the maximum changes in the index in patients with NASH, CKD and obesity, which testifies on the one hand, the presence of significant risk factors for the progression of atherosclerosis in these patients on the background of obesity, and on the other - on the favorable pathogenetic situation with regard to the progress of NASH. Thus, the development of NASH in patients with CKD and obesity is accompanied by a significant disorder of dyslipidemia with the highest among groups comparing with the increase in the content of cholesterol and low-density proatherogenic lipoprotein, a possible decrease in high-density anti-atherogenic lipoprotein and an increase in the atherogenicity index.

In patients of 1st and 2nd groups, a slight increase in cardiac glycemia was observed at 9.3% and 14.8%, respectively ( $p < 0.05$ ) compared with the control

group. Analysis of indicators of postprandial glycemia, obtained during Gamma glutamyltransferase (GGT), in patients of 1st and 2nd groups also showed an increase in glucose content in 120 min after loading - respectively by 16.5% and 31.2% ( $p < 0.05$ ) compared with indicators in the PHP group. Investigation of insulin content in blood on an empty stomach revealed hyperinsulinemia, which in patients of the 1st group exceeded the index in the PHP group by 1.9 times; in patients of the 2nd group insulin content exceeded the norm by 2.2 ( $p < 0.05$ ) times.

The results of the study showed that in patients with NASH, a significant increase in the content of NO in the blood was detected in comparison with the index in PHP ( $p < 0.05$ ) (Table 1) in group 1 - in 2.1 times, in the 2nd group - in 2.6 times ( $p < 0.05$ ). The role of nitrosative stress in the pathogenesis of NASH was proved, the confirmation of which is the increase in the concentration of nitrosothiols, peroxynitrite and other metabolites NO in the blood [2, 10]. Increased peroxynitrite formation due to the generation of NO by leukocytes is an important aspect of the damaging effect and inflammation process in NASH [3]. Pathological hyperproduction of NO by endothelial cells and leukocytes from inflammatory infiltrates in the liver contributes to the development of nitrosative stress in NASH. The established hypernitrate in blood may also be considered compensatory in response to hyperproduction of ET-1 in all observational groups. Thus, the content of ET-1 exceeded the index in PHP, respectively, in patients in the 1st group in 1.7 times, in the 2nd group - in 3.0 times ( $p_{1-2} < 0.05$ ). Confirmation of the presence of endothelial dysfunction (ED) in patients with NASH with CKD resulted in a probable growth of the number of desquamated endothelial cells (DEC) in the 2nd group of patients in 1.9 times ( $p_2 < 0.05$ ). Generation by neutrophils during the exacerbation of NASH of a significant number of active forms of oxygen and nitrogen and hyperproduction of endothelial cells and endometrial lymphocytes with progressive damage to the endothelium (growth of DEC) leads to significant ED, accompanied by mosaic angiospasm of the arteries due to hyperproduction of ET-1 and paretic vasodilatation of the veins of the portal vein system because of the hyperproduction of NO.

**Conclusions** Thus, in patients with non-alcoholic steatohepatitis on the background of obesity lipid distress syndrome with an increase in total cholesterol in blood, low density proatherogenic lipoproteins, and a deficiency in anti-atherogenic high-density lipoproteins is characterized. In a comorbid flow of non-alcoholic steatohepatitis and chronic kidney disease stage I and II on the background of obesity a deeper lipid imbalance (hypertriacylglycerolemia (2.1 times,  $p < 0.05$ ), hypercholesterolemia (1.5 times,  $p < 0.05$ ), including in the low density lipoprotein (1.8 times,  $p < 0.05$ ), decrease in the content of high density lipoprotein (1.8 times,  $p < 0.05$ ), increase in the atherogenic index (at 2, 7 times,  $p < 0.05$ ), as well as hyperproduction of the

endothelium of the NO and lymphocytes with progressive damage to the endothelium (growth of DEC)) was confirmed.

**The prospect of further scientific research in this direction** is the development of a method for the early prevention of non-alcoholic steatohepatitis on the background of obesity and the accompanying CKD of the 1st and 2nd stage.

#### References

1. Babak OYa, Kolesnikova EV, Syitnik KA. Profilakticheskie meropriyatiya pri nealkogolnoy zhirovoy bolezni pecheni: suschestvuet li sposob snizit risk razvitiya zabolevaniya? [Preventive measures for non-alcoholic fatty liver disease: is there a way to reduce the risk of the disease?]. *Suchasna gastroenterol.* 2013; 3 (71): 103-9. [Russian]
2. Bueverov AO, Bogomolov PO. Nealkogol'naya zhirovaya bolezni pecheni: obosnovanie patogeneticheskoy terapii [Non-alcoholic fatty liver disease: rationale for pathogenetic therapy]. *Klinicheskie perspektivy v gastroenterologii, gepatologii.* 2009; 1: 3-9. [Russian]
3. Anstee QM, Targher G, Day CP. Progression of NAFLD to diabetes mellitus, cardiovascular disease or cirrhosis. *Nat Rev Gastroenterol Hepatol.* 2013; 10: 330-44. PMID: 23507799. DOI: 10.1038/nrgastro.2013.41
4. Baumgarten M., Gehr T. Chronic kidney disease: detection and evaluation. *American Family Physician.* 2011; 84(10): 1138-48.
5. Brunt E.M., Kleiner D.E., Wilson L.A. et al. Nonalcoholic fatty liver disease (NAFLD) activity score and the histopathologic diagnosis in NAFLD: distinct clinicopathologic meanings. *Hepatology.* 2011;53(3):810-20.
6. Chalasani N., Younossi Z., Lavine J.E. et al. Diagnosis and Management of Non-alcoholic Fatty Liver Disease: Practice Guideline by the American Association for the Study of Liver Diseases, American College of Gastroenterology, and the American Gastroenterological Association. *Am. J. Gastroenterol.* 2012;107: 811-26.
7. Castera L, Vilgrain V, Angulo P. Noninvasive evaluation of NAFLD. *Nat Rev Gastroenterol Hepatol.* 2013; 10 (11): 666-75. PMID: 24061203. DOI: 10.1038/nrgastro.2013.175
8. Cohen E et al. A longitudinal assessment of the natural rate of decline in renal function with age. *J Nephrol.* 2014; 27(6):635-41.
9. Gambino R., Cassader M., Pagano G. Meta-analysis: natural history of non-alcoholic fatty liver disease (NAFLD) and diagnostic accuracy of non-invasive tests for liver disease severity. *Ann. Med.* 2011;43:617-49.
10. Nascimbeni F., Pais R., Bellentani S. et al. From NAFLD in clinical practice to answers from guidelines. *J. Hepatol.* 2013; 59(4):859-71.
11. Webb M., Yeshua H., Zelber-Sagi S. et al. Diagnostic value of a computerized hepatorenal index for sonographic quantification of liver steatosis. *Am. J. Roentgenol.* 2009; 192(4): 909-14.