

МІНІСТЕРСТВО ОХОРОНИ ЗДОРОВ'Я УКРАЇНИ
БУКОВИНСЬКИЙ ДЕРЖАВНИЙ МЕДИЧНИЙ УНІВЕРСИТЕТ



МАТЕРІАЛИ
106-ї підсумкової науково-практичної конференції
з міжнародною участю
професорсько-викладацького колективу
БУКОВИНСЬКОГО ДЕРЖАВНОГО МЕДИЧНОГО УНІВЕРСИТЕТУ
03, 05, 10 лютого 2025 року

Конференція внесена до Реєстру заходів безперервного професійного розвитку,
які проводитимуться у 2025 році №1005249

Чернівці – 2025

УДК 61(063)

М 34

Матеріали підсумкової 106-ї науково-практичної конференції з міжнародною участю професорсько-викладацького колективу Буковинського державного медичного університету (м. Чернівці, 03, 05, 10 лютого 2025 р.) – Чернівці: Медуніверситет, 2025. – 450 с. іл.

У збірнику представлені матеріали 106-ї науково-практичної конференції з міжнародною участю професорсько-викладацького колективу Буковинського державного медичного університету (м. Чернівці, 03, 05, 10 лютого 2025 р.) зі стилістикою та орфографією у авторській редакції. Публікації присвячені актуальним проблемам фундаментальної, теоретичної та клінічної медицини.

Загальна редакція: професор Геруш І.В., професорка Годованець О.І., професор Безрук В.В.

Наукові рецензенти:
професор Батіг В.М.
професор Білоокий В.В.
професор Булик Р.Є.
професор Давиденко І.С.
професор Дейнека С.Є.
професорка Денисенко О.І.
професор Заморський І.І.
професорка Колоскова О.К.
професорка Кравченко О.В.
професорка Пашковська Н.В.
професорка Ткачук С.С.
професорка Тодоріко Л.Д.
професорка Хухліна О.С.
професор Чорноус В.О.

ISBN 978-617-519-135-4

© Буковинський державний медичний
університет, 2025

Conclusions. Children with acute streptococcal tonsillopharyngitis exhibited a notably higher risk of cardiac performance decline with measured physical activity after one year.

Sazhyn S.I.

IS BODY MASS INDEX ASSOCIATED WITH POOR LEVEL OF ASTHMA CONTROL IN SCHOOL-AGE CHILDREN?

Department of Pediatrics and Children Infectious Diseases

Bukovinian State Medical University

Introduction. Asthma is a common, chronic respiratory disease affecting 1–29% of the population in different countries. According to Global Strategy for Asthma Management and Prevention (GINA) the long-term goal of asthma treatment is to achieve the best possible long-term outcomes for the patient. It includes symptom control and future risk of adverse outcomes. Poor symptom control is burdensome to patients and increases the risk of exacerbations, but patients with good symptom control can still have severe exacerbations. Obesity is known to be a risk factor for developing bronchial asthma in children, but there are difference opinions about the effect of overweight on asthma symptom control and future adverse outcomes.

The aim of the study. To evaluate the association between body mass index (BMI) and level of asthma control in school-age children to improve management of bronchial asthma.

Material and methods. On the base of the Chernivtsi Regional Children Clinical Hospital were observed 62 children who are suffered from bronchial asthma. According to the BMI of patients two clinical groups were formed. The first (I) group included 38 patients with BMI less than 20.99 kg/m^2 , the second (II) group was formed of 24 children with $\text{BMI} \geq 21.0 \text{ kg/m}^2$. The clinical groups were compared by gender, place of residence, severity and duration of disease. The «Asthma control test» (ACT) was used to determine the symptom control. Pulmonary functional tests (initial forced expiratory volume in the first second (FEV₁), peak expiratory flow (PEF) and airways reversibility) were performed to assess potential risk of future adverse outcomes like asthma attack. The impact of risk factors was evaluated by attributable risk, relative risk, odds ratio and their 95% confidence intervals (CI).

Results. According to the results of the ACT the average value of the scores was 20.2 (95% CI 19.3-21.2) in patients of I clinical group versus 17.1 (95% CI 16.0-18.5) ($P<0.05$) in school-age children with overweight. Thus, the attributable risk of not achieving control over asthma symptoms in children of the II clinical group was 28.2%, relative risk – 1.7 (95% CI 1.0-3.0) with odds ratio – 3.2 (95% CI 0.8-13.0). At the same time, there was no significant difference in spirometric indices in the comparison groups. The average value of the initial FEV₁ was 97.6% versus 96.2%, PEF – 82.6% versus 81.5%, bronchodilator responsiveness after inhalation of short-acting β_2 -agonist (a test with a result of 12.0% or higher is considered positive) – 7.2% and 9.6% among patients of I and II clinical group. Therefore, overweight in children does not have a significant effect on the risk of future adverse outcomes like asthma exacerbation.

Conclusions. Overweight in asthmatic children is associated with poorer symptom control but does not significantly affect the risk of future adverse outcomes. The lower level of asthma symptom control among children with overweight allows strongly recommend combination of low-dose inhalation corticosteroids and formoterol (track 1) as a reliever therapy.

Білоус Т.М.

ВМІСТ БІОМАРКЕРІВ РЕМОДЕЛІНГУ БРОНХІВ У МОКРОТИННІ ДІТЕЙ, ХВОРИХ НА БРОНХІАЛЬНУ АСТМУ, ЗА РІЗНОГО ЇЇ ДЕБЮТУ

Кафедра педіатрії та дитячих інфекційних хвороб

Буковинський державний медичний університет

Вступ. Процес персистування алергічного запалення в бронхах супроводжується їх морфологічними змінами у вигляді ремоделінгу та формуванням нечутливості до бронхорозширювальних препаратів. Виходячи з цього, представлялося доцільним вивчити біомаркери ремоделінгу дихальних шляхів у мокротинні хворих, що дозволило б у